



# IJPPR

INTERNATIONAL JOURNAL OF PHARMACY & PHARMACEUTICAL RESEARCH  
An official Publication of Human Journals

ISSN 2349-7203




Human Journals

**Research Article**

February 2018 Vol.:11, Issue:3


© All rights are reserved by Soumen Pattanayek et al.

## Formulation and Evaluation of Ketoprofen Loaded Nanoparticulate Gel for Topical Delivery



**IJPPR**  
INTERNATIONAL JOURNAL OF PHARMACY & PHARMACEUTICAL RESEARCH  
An official Publication of Human Journals

ISSN 2349-7203



**Soumen Pattanayek\*<sup>1</sup>, S.B.Puranik<sup>2</sup>**

<sup>1</sup>*Teva Pharmaceuticals 1776 Aspen Ln Weston, FL, 33327*

<sup>2</sup>*Drishti Institute of Distance Learning, Bangalore, Karnataka, India.*

**Submission:** 25 January 2018  
**Accepted:** 31 January 2018  
**Published:** 28 February 2018



HUMAN JOURNALS

[www.ijppr.humanjournals.com](http://www.ijppr.humanjournals.com)

**Keywords:** Nanoparticles, Carbopol, topical delivery, Ketoprofen

### ABSTRACT

The objective of the present study was to prepare ketoprofen loaded nanoparticles and to incorporate these nanoparticle single. Many drugs show poor permeation due to the barrier stratum corneum when delivered through the skin. To overcome this problem and to enhance the permeation of drug into the skin and also to reduce the skin irritation the drug was formulated in the form of nanoparticles using chitosan as a polymer and these drug loaded nanoparticles were incorporated in a gel for more effective delivery of a drug into the skin. The prepared drug-loaded nanoparticulate gel was evaluated for physical appearance, homogeneity, pH, drug content, viscosity, spreadability, extrudability and in vitro diffusion studies. All the nanoparticulate gel formulations were clear without any aggregates, particles, and fibers indicating excellent homogeneity of all the formulations. The pH of the formulations was closer to a neutral indicating absence of skin irritation and damage. The viscosity was directly proportional whereas extrudability and spreadability were inversely proportional to the concentration of carbopol employed. Formulation NG4 showed sustained release for 24 h.

## INTRODUCTION:

Topical drug delivery can be defined as the application of a medication to the skin or mucous membrane for local or systemic action. Delivery of drugs via skin is effective and avoids first-pass effect and metabolic degradation associated with oral administration [1]. Development of successful topical drug delivery system has been limited in scope due to the significant penetration barrier provided by the stratum corneum (SC), the topmost skin layer. To overcome this barrier, numerous active and passive penetration enhancement methods have been assessed [2-3]. Although active methods including iontophoresis, electroporation, and microneedles have shown some efficiency, more work is needed to establish for their safety band and cost-effectiveness. Passive methods include the use of chemical penetration enhancers, supersaturated systems, prodrugs, liposomes, microemulsions, and colloidal polymeric suspensions. This approach provides design flexibility (with formulation optimization) and the possibility of application over a larger area of skin compared to active methods [4-5]. Chemical penetration enhancers have been intensively investigated over the years, but the concentrations required for improved penetration often lead to irritation or sensitization. The use of vesicular delivery systems such as liposomes, transfersomes, ethosomes, and niosomes is still limited due to stability issues and insufficient understanding of the mechanism of penetration across intact skin. Additionally, lipid-based colloidal carriers have shown improved drug penetration through the skin, but their limited drug loading and the phase stability remain to be addressed, restricting their clinical applicability. Dispersion of polymer-based nanoparticles in hydrophilic gels can further improve drug delivery to the skin. The gel can aid in creating a uniform dispersion of the carriers in the matrix and increase the contact time and deposition of the carriers on the skin, resulting in enhanced skin penetration of the payload [6-7].

Ketoprofen, a potent NSAID is a preferential inhibitor of cyclooxygenase -2 and has analgesic and anti-inflammatory activity, widely used in the treatment of rheumatoid arthritis, osteoarthritis, and another joint disease. The poor aqueous solubility and wettability of ketoprofen lead to difficulty in formulating oral and topical formulation. Ketoprofen is classified in the Biopharmaceutics Classification Scheme as class II drug. Since dissolution is the rate-limiting step in drug absorption, the poor water solubility in oral forms of ketoprofen results in low bioavailability due to incomplete absorption. In addition to absorption difficulties, oral formulations of ketoprofen can cause gastric mucosal damage, which may

result in ulceration and bleeding. Therefore a better oral or topical formulation can be developed by increasing the water solubility of drugs. Topical application of NSAID on the inflamed site can offer the advantage of delivering a drug directly to the disease site and producing its local effect. This occurs by avoiding gastric irritation and also reduced adverse systemic effect [8-10].

In the present work, an attempt was made to enhance the absorption and provides sustained release of drug by incorporating the drug in nanoparticles and further in a gel for effective delivery of the drug into the skin.

## **MATERIALS AND METHODS:**

### **MATERIALS:**

Ketoconazole was a gift sample from Emcure Pharmaceuticals Ltd., Pune. Chitosan, Poloxamer 188 and Sodium tripolyphosphate were procured from Sigma Aldrich, USA. Acetic acid was purchased from SD fine chemicals, India. Deionized water was obtained from Millipore filtration system, USA.

### **METHODS:**

#### **Preparation of nanoparticles [11-13]:**

Nanoparticles were prepared by using chitosan (CS) as polymer and sodium tripolyphosphate (STPP) as the cross linking agent by ionic gelation method. Initially, 0.4 % of chitosan solution in 1% acetic acid was prepared and pH was adjusted to 4. To the chitosan solution, drug solution was added slowly under constant stirring. 1.5 % w/v poloxamer 188 was added as the stabilizer. After 15 mins, 0.25% w/v STPP solution was added drop by drop slowly under magnetic stirring (Remi, India) at the constant speed of 1500 rpm for 2h and finally subjected to high-speed homogenizer (SB1, Thomas Scientific, USA) at 25,000 rpm for 1 h. The suspension of nanoparticles was then centrifuged at 10,000 rpm at 4 °C for 20 min and the separated nanoparticles were resuspended in purified water and then freeze-dried (Bioline Technologies, Thane, India) using 5% mannitol as cryoprotectant.

**Preparation of nanoparticulate gel [14]:**

The nanoparticulate gel was prepared by dissolving the required quantity of carbopol 934 in 100 ml distilled water containing 0.3 ml methylparaben (0.5% w/v) and stirred until it forms a clear transparent gel. Triethanolamine was slowly added and the pH was adjusted to 7. It was then allowed to stand for 24 h at room temperature. Then finally drug-loaded nanoparticles were incorporated in a gel by slow stirring. The various formulations prepared are shown in table 1.

**Table 1: Formulations of nanoparticulate Topical gel**

Ingredients	NG1	NG2	NG3	NG4	NG5	NG6
Ketoprofen (%w/w)	1	1	1	1	1	1
Carbopol 934 (%w/w)	0.5	1.0	1.5	2.0	2.5	3.0
Methylparaben (ml)	0.3	0.3	0.3	0.3	0.3	0.3
Triethanolamine (ml)	q.s	q.s	q.s	q.s	q.s	q.s
Water (ml) Up to	100	100	100	100	100	100

**Physical appearance and homogeneity [15]**

All the prepared nanoparticulate gels were tested for homogeneity by visual inspection after the gels have been set in the container. They were tested for their appearance and presence of any aggregates, particles, and fibers.

**Determination of pH**

The pH measurements were done using a digital pH meter (Thermo scientific) which was calibrated with buffer and the pH of the formulations was measured by dipping the glass electrode into the topical gel.

**Determination of drug content**

1.0 gm of the nanoparticulate gel was transferred to 50 ml volumetric flask and was diluted with ethanol. 5 ml of this solution was further diluted to 25 ml with ethanol. The drug content

was determined by measuring the absorbance at 254 nm using UV- Visible spectrophotometer.

#### **Determination of viscosity [15]**

The viscosity of the prepared formulations was determined at ambient temperature using Brooke field digital viscometer (DV-II + Pro) with spindle no. 6 at 5 rpm at room temperature and values were noted.

#### **Determination of spreadability [16]**

A weighed quantity (500 mg) of topical gel or gels was placed within a circle of 1 cm diameter pre-marked on a glass plate (10 X 10 cm). Another glass plate (10 X 10 cm) was placed on the gel. A weight of 500 g was allowed to rest on the upper glass plate for 5 min. The increase in the diameter due to gel spreading was noted.

#### **Determination of Extrudability**

The quantity ( $\text{g}/\text{cm}^2$ ) of topical gel extruded from lacquered aluminium collapsible tube on an application of weight (in grams) required to extrude at least 0.5 cm ribbon of topical gel in 10 seconds. The measurement of extrudability of each formulation was done in triplicates using the below equation and the average values are presented.

$$\text{Extrudability} = \text{Applied weight (in g)}/\text{Area (in cm}^2\text{)}$$

#### ***In vitro* diffusion studies through dialysis membrane [17]**

The membrane utilized was dialysis membrane number 70, having average flat width 29.31 mm, an average diameter of 17.5 mm and a capacity of 2.41 ml/cm (Molecular weight cut off 12000). The dialysis membrane was cut to size and boiled in distilled water for 1 hour. Then soaked in absolute alcohol for 1 hour and was finally soaked in pH 7.4 phosphate buffer saline for 24 hours. *In vitro*, drug release studies were carried out by taking required quantity of nanoparticulate gel on the dialysis membrane, which was mounted on the Franz diffusion cell, the top of which was clamped securely. The receptor compartment with pH 7.4 phosphate buffer saline was maintained at constant temperature of 37° C by circulating water bath. Samples were withdrawn from the receptor compartment at predetermined time

intervals and replaced with an equal volume of fresh buffer solution. The samples were analyzed for drug content using UV spectrophotometer at 254 nm.

## RESULTS AND DISCUSSION:

### Preparation of nanoparticulate gel:

Drug-loaded chitosan nanoparticles were prepared using ionotropic gelation method. In nanoparticle preparation, chitosan was the polymer and cross-linked with sodium tripolyphosphate, the cross-linker, by ionic linkage. In the preparation of gel, carbopol was used as gelling agent. One of the main ingredients of the formulation is the gelling agent. The concentration of viscosity enhancer or gel former is of immense value as a less concentration will lead to simple solution or lotion with very low consistency, while high concentration may lead to the formation of gels with high viscosity leading to the non-uniform distribution of drug and problem with the handling of gel.

### Physical appearance and homogeneity

All the nanoparticulate gel formulations were clear without any aggregates, particles, and fibers indicating excellent homogeneity of all the formulations. The results are indicated in Table 2. At low concentrations of carbopol (NG1 to NG3) a very thin gel that liquefies within 24h of preparation was formed. A gel containing 2.0% of carbopol 934 (NG4) formed the uniform and smooth gel that does not liquefy upon keeping. At further higher concentrations of the gelling agent (NG5 and NG6) the gel was very thick and stickier that could not be properly spread out.

**Table 2: Results of Physical appearance and homogeneity for gels**

Formulation code	Appearance	Homogeneity
NG1	Clear, white fluid	+++
NG2	Clear, white fluid	+++
NG3	Clear, white fluid	+++
NG4	Clear, white fluid	+++
NG5	Clear, thick white	+++
NG6	Clear, thick white	+++

Excellent+++ , Good++ , Satisfactory+

### Determination of pH

The pH of all the formulations was in the range of  $6.78 \pm 0.04$ -  $7.25 \pm 0.06$  (Table 3) enduring them most acceptable to avoid the risk of irritation upon application to the skin. This also indicated that the selected ingredients of the formulation did not alter the pH of the formulation.


**Table 2: Results for determination of pH for gels**

Formulation code	pH (mean $\pm$ SD*)
NG1	$6.72 \pm 0.04$
NG2	$6.78 \pm 0.03$
NG3	$6.82 \pm 0.08$
NG4	$6.95 \pm 0.05$
NG5	$7.25 \pm 0.06$
NG6	$7.15 \pm 0.04$

\*Standard deviation, n=3

### Determination of drug content


The drug content for all the prepared nanoparticulate gel formulations was found between 95.5% to 97.0%. The drug content analysis showed that the drug-loaded nanoparticles were uniformly and properly distributed in the gel formulations. The results obtained are shown in figure 1.



**Fig. 1: Drug content for the various formulation of gels (mean  $\pm$  standard deviation, n=3)**

### Determination of viscosity 52

Viscosity depends upon the type of polymer and also the concentration of polymer used. In general, viscosity increases with increasing the concentration of polymer.



**Fig. 2: Results of viscosity studies for the various formulation of gels (mean  $\pm$  standard deviation, n=3)**

### Determination of spreadability 53

Spreadability plays an important role in patient compliance and helps in the uniform application of gel to the skin as gels should spread easily upon application. Spreadability mainly depends on the viscosity of the gel. If the viscosity of the formulation was more then it affects the spreadability. Increase in viscosity leads to decrease in spreadability. Topical gels prepared with the low concentration of carbopol (NG1 to NG3) belonged to fluid gel category, having more spreadability values. The formulations prepared with the higher concentration of carbopol belonged to a stiff and semi stiff category. Increase in concentration of gelling agent in formulation resulted in the decrease of spreadability values (Table 3). The results indicated that the formulation can be applied easily without being the runoff. This assures that the formulation maintains a good wet contact time when applied to the site of application.



**Table 3: Results of determination of spreadability and extrudability for gels**

Formulation code	Spreadability (cm) (mean±SD*)	Extrudability (g/cm <sup>2</sup> ) (mean±SD*)
NG1	15.4±0.75	22.30±1.02
NG2	11.56±0.63	21.52±0.85
NG3	10.75±0.54	19.23±0.75
NG4	9.26±0.23	15.42±0.62
NG5	8.46±0.42	8.62±0.45
NG6	6.23±0.16	6.45±0.32


\*Standard deviation, n=3

### **Determination of Extrudability**

It was found that extrudability of the topical gel was a function of a concentration of carbopol. Extrudability was decreased with increase in the concentration of carbopol (table 3).

### ***In vitro* diffusion studies through the dialysis membrane**

Diffusion studies were carried out using Franz type diffusion cell for NG1 to NG6 formulations in pH 7.4 Phosphate buffer saline solution (Figure 3). The drug release from NG5 and NG6 was less due to more concentration of carbopol. As more concentration of carbopol was employed the drug was not able to completely diffuse through the thick gel barrier. Drug release was in the following order, NG6>NG5>NG4>NG3>NG2>NG1. The drug release profile of NG4 was greater. Hence NG4 can be considered as the optimized formulation.



**Fig.3: In vitro release of ketoprofen from formulations NG1 - NG6 (mean  $\pm$  standard deviation, n=3)**

**CONCLUSION:**

The aim of the present work was to formulate, develop and evaluate a topical gel containing ketoprofen. Topical gels have emerged as a promising drug delivery system for the delivery of hydrophobic drugs. Formulations were prepared and evaluated for physical parameters, drug content, viscosity, spreadability, extrudability and diffusion studies. Carbopol, a gelling agent has a direct influence on appearance, viscosity, spreadability and extrudability. As the concentration of gelling agent increases the viscosity of the formulation increased and release was retarded. From the results of the experimental work carried out, it can be concluded that the ketoprofen loaded nanoparticles incorporated in gels can be employed for effective sustained topical delivery of lipophilic drugs and drugs with a high molecular weight which shows poor permeation into the skin.

**REFERENCES:**

1. Barkin RL. Topical Nonsteroidal Anti-Inflammatory Drugs: The Importance of Drug, Delivery, and Therapeutic Outcome. *Am J Ther.* 2015 Sep-Oct;22(5):388-407.
2. Stanos SP, Galluzzi KE. Topical therapies in the management of chronic pain. *Postgrad Med.* 2013 Jul;125(4 Suppl 1):25-33.
3. Rehman K, Zulfakar MH. Recent advances in gel technologies for topical and transdermal drug delivery. *Drug Dev Ind Pharm.* 2014 Apr;40(4):433-40.
4. Singla SK, Sachdeva V. Current and emerging lipid-based systems for transdermal drug delivery. *Ther Deliv.* 2015; 6(9):1063-70.
5. Donnelly RF, Garland MJ, Alkilani AZ. Microneedle-iontophoresis combinations for enhanced transdermal drug delivery. *Methods Mol Biol.* 2014; 1141:121-32.
6. Vadlamudi HC, Narendran H, Nagaswaram T, Yaga G, Thanniru J, Yalavarthi PR. Microemulsions based transdermal drug delivery systems. *Curr Drug Discov Technol.* 2014;11(3):169-80.

7. Jain A, Jain P, Kurmi J, Jain D, Jain R, Chandel S, Sahu A, Mody N, Upadhaya S, Jain A. Novel strategies for effective transdermal drug delivery: a review. *Crit Rev Ther Drug Carrier Syst.* 2014;31(3):219-72.
8. Lauterbach A, Müller-Goymann CC. Applications and limitations of lipid nanoparticles in dermal and transdermal drug delivery via the follicular route. *Eur J Pharm Biopharm.* 2015 Nov;97(Pt A):152-63.
9. Ferreira LM, Cervi VF, Gehrcke M, da Silveira EF, Azambuja JH, Braganhol E, Sari MH, Zborowski VA, Nogueira CW, Cruz L. Ketoprofen-loaded pomegranate seed oil nanoemulsion stabilized by pullulan: Selective antiangioma formulation for intravenous administration. *Colloids Surf B Biointerfaces.* 2015 Jun 1;130:272-7.
10. Abd-Elrahman AA, El Nabarawi MA, Hassan DH, Taha AA. Ketoprofen mesoporous silica nanoparticles SBA-15 hard gelatin capsules: preparation and *in-vitro/in-vivo* characterization. *Drug Deliv.* 2016 Nov;23(9):3387-3398.
11. Kawashima Y, Handa T, Kasai A, Takenaka H, Lin SY, Ando Y. A novel method for the preparation of controlled-release theophylline granules coated with a polyelectrolyte complex of sodium polyphosphate chitosan. *J Pharm Sci.* 1985;74(3):264-8.
12. Kleine-Brueggeney H, Zorzi GK, Fecker T, El Gueddari NE, Moerschbacher BM, Goycoolea FM. A rational approach towards the design of chitosan-based nanoparticles obtained by ionotropic gelation. *Colloids Surf B Biointerfaces.* 2015 Nov 1;135:99-108
13. Jing ZW, Jia YY, Wan N, Luo M, Huan ML, Kang TB, Zhou SY, Zhang BL. Design and evaluation of novel pH-sensitive ureido-conjugated chitosan/TPP nanoparticles targeted to *Helicobacter pylori*. *Biomaterials.* 2016 Apr;84:276-85.
14. Nornoo AO, Wulz J, Yoon H, Nan Y, Lese M. Impact of the chemical and physical stability of ketoprofen compounded in various pharmaceutical bases on its topical and transdermal delivery. *Pharm Dev Technol.* 2016 Mar;21(2):204-13.
15. Sakeena MH, Muthanna FA, Ghassan ZA, Kanakal MM, Elrashid SM, Munavvar AS, Azmin MN. Formulation and in vitro evaluation of ketoprofen in palm oil esters nanoemulsion for topical delivery. *J Oleo Sci.* 2010; 59(4):223-8.
16. Nagai N, Iwamae A, Tanimoto S, Yoshioka C, Ito Y. Pharmacokinetics and Antiinflammatory Effect of a Novel Gel System Containing Ketoprofen Solid Nanoparticles. *Biol Pharm Bull.* 2015;38(12):1918-24.
17. Richards H, Thomas CP, Bowen JL, Heard CM. *In-vitro* transcutaneous delivery of ketoprofen and polyunsaturated fatty acids from a pluronic lecithin organogel vehicle containing fish oil. *J Pharm Pharmacol.* 2006 Jul;58(7):903-8.