



IJPPR

INTERNATIONAL JOURNAL OF PHARMACY & PHARMACEUTICAL RESEARCH
An official Publication of Human Journals

ISSN 2349-7203





Human Journals

Review Article

March 2019 Vol.:14, Issue:4

© All rights are reserved by Dinesh Kumar Meena

In-Vivo Models Used for Pre-Clinical Evaluation of Anti-Ulcer Activity: A Review

	
<p>Dinesh Kumar Meena^{*1}</p> <p><i>¹Department of Pharmacology, Jawaharlal Institute of Postgraduate Medical Education and Research (JIPMER), Puducherry-605006, India.</i></p> <p>Submission: 23 February 2019 Accepted: 28 February 2019 Published: 30 March 2019</p>	



www.ijppr.humanjournals.com

Keywords: Gastric ulcer, Ulcer score, Ulcer index, Percentage protection. Percentage inhibition, Non-steroidal anti-inflammatory drugs

ABSTRACT

Gastric ulcer disease has become a disease predominantly affecting the older population, with the peak incidence occurring between 55 and 65 years of age. It can occur in any part of gastrointestinal tract. Many medications are available for management of gastric ulcer. Prolonged use of these drugs may lead to serious adverse effects. Advanced in the discovery of more effective and safe anti-ulcer agent is due to the introduction of large number of newer experiment methods to evaluate their anti-ulcer activity in different types of gastric ulcers. Several *in-vivo* models of gastric damage have been characterized and are primary tools to identify the anti-ulcer property of many new and existing drugs.

INTRODUCTION:

Ulcers are lesions of the skin or mucous membrane characterized by the superficial inflamed dead tissue¹. Peptic ulcer is the most predominant gastrointestinal disease². Studies showed that gastric ulcer occurs at least 10% of the world population⁴. Peptic ulcer caused by a lack of balance between the gastric aggressive factors and gastric protective factors^{3,5}. Aggressive factors include increased secretion of HCL and pepsin, inadequate dietary habits, free oxygen radicals, consumption of NSAIDs and alcohol, stress and infection of *Helicobacter pylori*. Gastric protective factors include adequate gastric blood flow, secretion of prostaglandins, mucous, nitric oxide, bicarbonates and growth factors^{6,7}.

Drugs such as anticholinergic, histamine H₂ receptor antagonists, antacids and proton pump inhibitors are commonly used for treatment of peptic ulcer⁸. Prolong use of these drugs may lead to serious adverse effects like thrombocytopenia, nephrotoxicity, hepatotoxicity and impotence^{8,9}. Due to unpleasant side effects of existing anti-ulcer drugs, there is need of more effective and safe treatment for ulcers.

There are several models used to evaluate anti-ulcer activity of existing as well as new drugs. This review mainly focus on various *in-vivo* models available for pre-clinical evaluation of anti-ulcer activity of drugs.

***In-Vivo* Models Used For Pre-Clinical Evaluation of Anti-Ulcer Activity**

1. Water-immersion stress or cold-restraint induced gastric ulcer model
2. Non steroid anti- inflammatory drugs (NSAIDs) induced gastric ulcer model
3. Ethanol induced gastric ulcer model
4. Acetic acid induced gastric ulcer model
5. Histamine induced gastric ulcer model
6. Reserpine induced gastric ulcer model
7. Serotonin induced gastric ulcer model
8. Pylorus ligated induced ulcer model

9. Diethyl dithiocarbonate (DDC) induced ulcer model
10. Methylene blue induced ulcer model
11. Ischemia-reperfusion (I-R) induced gastric ulcer model
12. Cysteamine-induced duodenal ulcer model

1. Water-immersion stress or cold-restraint induced gastric ulcer model

Principle: In this model, gastric ulcers are induced by water-immersion stress or cold restraint stress in rats or mice. Stress induces ulcers by release of histamine which leads to an increased secretion of gastric acid, reduction in mucous production, reflux of pancreatic juice, and impairs gastric blood flow and increased gastro-intestinal motility^{10,11,12,13}.

Procedure:

Water-immersion stress induced ulcer model¹⁴:

- Animals are fasted for a period of 24-36 hours prior to the experiment.
- Animals treated with vehicle or test drug or reference drug.
- 30 minutes later, animals are placed individually in each compartment of a stress cage and immersed vertically up to xiphoid level in water bath and kept for 7 hours which result in induction of ulcers.
- 7 hours later, animals are sacrificed, stomach is dissected out and severity of ulcers is examined by calculating ulcer index.

Cold-restraint stress induced ulcer model^{14,15}: Cold water immersion accelerates the development of ulcers in restraint animals. Wister rates are used for experiment.

- Animals are fasted for 16 hours prior to the experiments.
- Test compound is administered orally.
- 1 hour later, animals are individually restraint in restraint cages vertically for 2 hours.
- Animals are immersed in water at 22°C for 1 hour.

- Evans's blue, in dose of 30 mg/kg is injected intravenously via the tail vein.
- Animals are sacrificed 10 minutes later.
- Stomach is removed and ligate at both ends.
- Stomach is filled with saline and kept overnight.
- On the next day, stomach is opened along the greater curvature, washed in warm water and examined for ulcer lesions.

2. NSAIDs induced gastric ulcer model: NSAIDs like aspirin, indomethacin and ibuprofen are the second most common cause of gastric ulcer¹⁶.

Principle: NSAIDs cause ulcers by inhibiting prostaglandins synthesis by inhibiting cyclooxygenase enzyme in COX pathway. Prostaglandins play a protective role via stimulating the secretion of bicarbonates and mucous, maintaining blood flow and regulating mucous cell turn over and repair^{17,18}.

Procedure¹⁶:

- Animals are fasted for 24-36 hours.
- NSAID (aspirin or indomethacin) using appropriate vehicle (water or 1% carboxymethylcellulose) is orally administered.
- After 1 hour, animals are treated with test drug.
- 4 hours later, animals are sacrificed, stomach is removed and severity of ulcer is measured.
- Dosage of NSAIDs used to induce ulcers are as following:

Aspirin: 150 mg/kg of body weight.

Indomethacin: 40-100 mg/kg of body weight

3. Ethanol-induced gastric ulcer model

Principle: Ethanol causes ulcer lesions by exposing the gastric mucous to the hydrolytic and photolytic actions of HCL and pepsin^{19,20}.

Procedure^{15,16}: Wister rats are used for experiment.

- Animals are fasted for 18 hours.
- Test drug is given to animals orally.
- 30 minutes later, 1 ml of absolute ethanol is administered orally.
- 1 hour later, animals are sacrificed and their stomachs are dissected out.
- Stomach are opened along the greater curvature, washed with warm water and examined for ulcer severity.

4. Acetic acid induced gastric ulcer model

This method used for chronic peptic ulcers. This method is suitable to evaluate the effect of potential drugs and also to test the drug on the healing of chronic ulcers. Method can also use to screen ant secretory and ulcer protective effect of drugs^{16,21}.

Procedure^{15,16}: albino rats are used for the experiment.

- Animals are fasted for 24-36 hours.
- Animals are anaesthetised.
- A flexible plastic catheter with an outside diameter of 2 mm is inserted up to 8 cm in colon via anus, through which 2 ml of diluted acid (4%) is introduced into colon.
- Animals are then kept into head down position for 2 minutes to prevent leakage of the acetic acid solution.
- After 24 hours, animals are sacrificed, stomach are removed and opened with greater curvature.
- Ulcer index is calculated to examine ulcer severity.

5. Histamine-induced gastric ulcer model

Principle: Histamine released from mast cells and binds with receptors present on the surface of parietal cells which leads to activation of adenylyl cyclase. This adenylyl cyclase converts ATP into c-AMP. This conversion enhances secretion of HCL from parietal cells²².

Procedure^{15,16}: Male guinea pigs are used for the experiment.

- Animals are fasted for 36 hours.
- Histamine acid sulphate in dose of 50 mg is injected intraperitoneally.
- To prevent histamine toxicity, promethazine hydrochloride in dose of 5 mg is injected intraperitoneally 15 minutes before and 15 minutes after the histamine injection.
- Test drug is administered 30-45 minutes later of histamine injection.
- After 4 hours, animals are sacrificed, stomach removed and dissected.
- Ulcer index is calculated to examine the severity of ulcers.

6. Reserpine-induced gastric ulcer model

Principle: Reserpine acts on cholinergic system. Reserpine increases histamine secretion by causing degranulation of gastric mast cells²³.

Procedure^{15,16}: Female Sprague - Dawley rats are used for the experiment.

- Animals are fasted for 48 hours.
- Test drug is administered intraperitoneally.
- Half an hour later, reserpine in dose of 15 mg/kg is administered intraperitoneally.
- 4 hours later, animals are sacrificed, stomach are removed and dissected.
- Ulcer index is calculated.

7. Serotonin-induced gastric ulcer model

Principle²⁴: Serotonin acts as vasoconstrictor which reduces gastric mucosal blood flow and leads to acute mucosal injury.

Procedure¹⁶:

- Animals are fasted for 24 hours.
- Serotonin creatin sulfate (0.5 ml of 20-50 mg/kg body weight) is administered subcutaneously.
- Animals are sacrificed after 6 hours; stomach is dissected out and examined for ulcer severity.

8. Pylorus ligated induced ulcer model

Principle¹⁶: The ligation of the pyloric end of the stomach causes accumulation of gastric acid in the stomach which leads to development of ulcers.

Procedure^{15,16}: Wister rats (150 to 180 gm weight) are used for the experiment.

- Animals are fasted for 48 hours.
- Animals are anaesthetized and a 1-inch midline abdominal incision is given below the xiphoid process.
- The pylorus is carefully lifted out and ligated without damaging its blood supply.
- The stomach is now replaced and the abdominal wall closed with sutures.
- The test compound is administered either orally or subcutaneously.
- 10-19 hours later, animals are sacrificed and stomachs are dissected out.
- Contents of the stomach are drained into a graduated centrifuge tube and their acidity determined by titration with 0.1 N NaOH.
- Stomach is opened along its greater curvature and ulcer index is calculated.

9. Diethyl Dithiocarbonate (DDC) induced ulcer model

Principle²⁵: DDC induces ulcers through the mobilization of superoxide and hydroxyl radicals. Superoxide radicals and hydroxyl radicals plays a pathogenic role in development of ulcers.

Procedure^{14,26}: This model is used to assess the antioxidant activity and cytoprotective activity of drug.

- Animals are fasted for 24 hours.
- Acute glandular lesions are induced by subcutaneous injection of 1 ml of DDC in saline followed by oral dose of 1 ml of 0.1 N HCl.

10. Methylene Blue induced ulcer model

Principle^{27,28}: Methylene blue is a synthetic drug. It is known to generate superoxide radical ions by uncoupling of ATPase. In addition, it also have anticholinergic activity. Drugs with c activity and proton pump inhibitory activity can be assessed by using this model.

Procedure¹⁶:

- Animals are fasted for 24 hours.
- Methylene blue is administered at a dose of 125 mg/kg of body weight orally followed by the administration of test drug.
- Animals are sacrificed after 4 hours of methylene blue administration.
- Stomachs are dissected out and ulcer index is calculated.

11. Ischemia-Reperfusion (I-R) induced gastric ulcer model.

Principle²⁹: Reperfusion of gastro intestine following ischemia leads to formation of free radicals which results in development of erosion and ulceration in the gastric mucosa.

Procedure³⁰:

- Animals are fasted for 24 hours.

- Animals are anesthetized.
- Laparotomy is performed and esophageal and pyloric ends of the stomach are clamped using bull god clips.
- Celiac artery is then clamped at a point of 0.5 cm distal from the branch to the aorta for 30 min.
- GI is then reperfused for 20 min.
- Animals are sacrificed, stomachs are dissected out and ulcer index is calculated.

12. Cystamine induced duodenal ulcer model

Principle^{31,32}: Cystamine develops formation of duodenal ulcers by stimulating gastric acid secretion and inhibiting the secretion of alkaline mucous from Brunner's gland.

Procedure³³: There are two types of duodenal ulcers i.e. acute and chronic.

- Acute ulcers can be produced by administering single dose (400 mg/kg of body weight) of cystamine HCL.
- Chronic ulcers can be produced by administering (400mg/kg of body weight) of cystamine HCL twice at an interval of 4 hours.
- Cut opened along the antimesenteric side and ulcer areas are measured.

Parameters To Be Calculated³⁴

There are three parameters i.e ulcer index, % protection ratio and % curative ratio, to be calculated by using method described by Tokagi and Okabe to evaluate anti-ulcer activity of drug in in-vivo models.

Steps:

1. Give score based on ulcer severity.
2. Calculate Ulcer Index (UI) based on ulcer score.
3. % protection ratio and % curative ratio can be calculated by using Ulcer Index. (UI).

1. Scoring of ulcers based on ulcer severity:

Score	Ulcer severity
0	No lesions
1	mucosal oedema
2	1-5 small lesions (1-2 mm in size)
3	> 5 small or intermediate (3-4 mm in size) lesions
4	≥ 2 intermediate lesions or 1 gross (> 4 mm in size) lesion
5	Perforated lesions

2. Calculation of Ulcer Index (UI) based on ulcer score: By using ulcer score as described above, ulcer index can be calculated as following:

$$\text{Ulcer Index (UI)} = \frac{\text{Total ulcer score}}{\text{Number of animals ulcerated}}$$

3. Calculation of % protection ratio and % curative ratio by using Ulcer Index.

% protection ratio =

$$\frac{\text{UI of ulcerogen treated group}}{\text{UI of ulcerogen treated}} - \frac{\text{UI of drug pre-treated group}}{\text{UI of ulcerogen treated}}$$

% curative ratio =

$$\frac{\text{UI of ulcerogen treated group}}{\text{UI of ulcerogen treated}} - \frac{\text{UI of drug treated group}}{\text{UI of ulcerogen treated}}$$

DISCUSSION

Peptic ulcer is most common gastrointestinal disease that mainly caused by *H. pylori* infection and high intake of NSAIDs. Pre-clinical evaluation of new or existing antiulcer drug can be performed by using appropriate *in-vivo* models. Several models are developed to evaluate antiulcer activity of natural as well as synthetic drugs. *In-vivo* models may also be used to assess any toxic effects of test drug.

REFERENCES

1. Chan FKL, Graham DY. Review article: prevention of non-steroidal anti-inflammatory drug gastrointestinal complications - review and recommendations based on risk management. *Aliment Pharmacolo Ther.* 2004;19:1051-1061.
2. R.K. Goyal. Elements of Pharmacology, B.S. Shah Prakashan, New Delhi, India, 17th edition, 2008.
3. P. Malfertheiner, F.K. Chan, and K.E. McColl. Peptic ulcer disease. *The Lancet.* 2009;347(9699):1449-1461.
4. Shimoyama AT, Santin JR, Machado ID, de Oliveria e Silva AM, de Melo IL, Mancini-Filho J, et al. Antiulcerogenic activity of chlorogenic acid in different models of gastric ulcer. *Naunyn-Schiedebergs Arch Pharmacol.* 2013;386:5-14.
5. C.V. Rao, K. Sairam, and R.K. Goel. Experimental evaluation of Bopocamonniera on rat gastric ulceration and secretion. *Indian Journal of Physiology and Pharmacology.* 2000;44(4):435-441.
6. Sowndhararajan K, Kang SC. Protective effect of ethyl acetate fraction of *Acasia Ferruginea* DC. Against ethanol-induced gastric ulcer in rats. *J Ethnopharmacol.* 2013;148:175-181.
7. Lemos M, Santin JR, Mizuno CS, Boeing T, de Sousa JPB, Nanayakkara D, et al. Copaiteralangsdorffii: Evaluation of potential gastroprotective of extract and isolated compounds obtained from leaves. *Rev Bras Formacogn.* 2015;25:238-245.
8. Chan FK, leung WK, Peptic-ulcer disease. *The Lancet.* 2002;360:933-941.
9. Sheen FK. Triadafilopulos G. Adverse effect of long term proton pump inhibitor pathway. *Dig Dis Sci.* 2011;56:931-950.
10. H. Kitagawa, M. Fujiwara, and Y. Osumi. Effects of water-immersion stress on gastric secretion and mucous blood flow in rats. *Gastroenterology.* 1979;77(2):298-302.
11. P. H. Guth. Gastric blood flow in restraint stress. *The American Journal of Digestive Diseases.* 1972;17(9):807-813.
12. M. N. Peters and C. T. Richardson. Stressful life event, acid hypersecretion and ulcer disease. *Gastroenterology.* 1983;84(1):114-119.
13. D. A. Brodie and H. M. Hanson. A study of the factors involved in the production of gastric ulcers by the restraint techniques. 1960;38:353-360.
14. Thabrew MI and Mrawwawala LDAM. An overview of In vivo and In vitro Models that can be used for evaluating Anti-Gastric Ulcer Potential of Medicinal Plants. *Austin Biol.* 2016;1(2):1007.
15. Michael Buenor Adinortey et al. In vivo Models used for Evaluation of Potential Antigastroduodenal Ulcer Agents. *Hindawi Publishing Corporation.* 2013;1-12.
16. K. D. Rainsford. The effect of 5-lipoxygenase inhibitors and leukotriene antagonists on the development of gastric lesions induced by nonsteroidal anti-inflammatory drugs in the mice. *Agents and Actions.* 1987;21(3,4):316-319.
17. J. Hayliar and I. Bjarnason. NSAIDs, COX-2 inhibitors, and the gut. *The Lancet.* 1995;346(8974):521-522.
18. P. J. Oates and J. P. Hakkinen. Studies on the mechanism of ethanol-induced gastric damage in rat. 1988;94(1):10-21.
19. G. Sener, K. Pasakalogu, and G. Ayanoglu-Dulger. Protective effect of increasing doses of famotidine, omeprazole, lansoprazole, and melatonin against ethanol-induced gastric damage in rats. *Indian Journal of Pharmacology.* 2004;36(3):171-174.

20. K. Takagi, S. Okabe, and R. Saziki. A New method for the production of chronic gastric ulcers in rats and the effect of several drugs on its healing. *Japanese Journal of Pharmacology*.1970;19(3):418-421.
21. L. J. Hay, R. L. Varco, C. F. Code, and O. F. Wangsteen. Experimental production of gastric and duodenal ulcers in laboratory animals by intramuscular injection of histamine in beeswax. *The Journal of Surgery, Gynecology and Obstetrics*.1942;74:70-182.
22. S. Singh. Evaluation of gastric anti-ulcer activity of fixed oil of *ocimumbasilicum* Linn. and its possible mechanism of action. *Indian Journal of Experimental Biology*.1999;37(3):253-257.
23. K. J. Lepard and R. L. Stephens. Serotonin inhibits gastric acid secretion through a 5-hydroxy tryptamine like receptor in the rat. *Journal of Pharmacology and Experimental Therapeutics*.1994;270(3):1139-1147.
24. A. S. Salim. Protection against stress-induced acute gastric mucosal injury by free radicals scavengers. *Intensive Care Medicine*.1991;17(8):455-460.
25. S. Oka, Oginok, I. Hobara et al. Role of reactive oxygen species in diethyldithiocarbamate induced gastric ulcer in the rat. *Experimentia*.1990;46:281-283.
26. D.I. Shah, D. D. Santani, and S. S. Goswami. A novel use of methylene blue as a pharmacological tool. *Journal of Pharmacology and Toxicology Methods*.2006;54(3):273-271.
27. M. Pfaffendorf, T. A. Bruning, H. D. Batink, and P.A. Van Zwieten. The interaction between methylene blue and the cholinergic system. *British Journal of Pharmacology*.1997;122(1):95-98.
28. V. Kumar, A.K. Abbas, and N. Fausto. Pathological basis of disease. In Robbins and Cotran. New Delhi Saunders, New Delhi, India, 7th edition.2003.787-802.
29. A. Onen, Z. Kanay, C. Guzel, D. Kurt, and K. Ceylon. The effect of allopurinol on stomach mucosal barrier of rats subjected to ischemia reperfusion. *Turkish Journal of Medical Sciences*.2000;30(5):449-452.
30. Y. Ishii, Y. Fujii, and M. Homma. Gastric acid stimulating action of cysteamine in the rats. *European Journal of Pharmacology*. 1976;36(2):331-336.
31. H. Tamaki, Y. Onoda, and T. Kashida. Gastric secretion and duodenal ulcer formation induced by cysteamine in rats. *Japanese Journal of Pharmacology*.1978;28(4):647-649.
32. H. Selye and S. Szabo. Experimental model for production of perforating duodenal ulcers by cysteamine in the rat. *Nature*;244(5416):458-459.
33. P. S. Olsen, P. Kirkegaard, J. Christiansen, and S.S. Paulsen. Healing of acute and chronic experimental ulcer in rats. *Scandinavian Journal of Gastroenterology*.1982;17(78):1250.
34. Jyoti Gupta | Dinesh Kumar, Ankit Gupta. Evaluation of gastric anti-ulcer activity of methanolic extract of *Cayratia trifolia* in experimental animals. *Asian Pacific Journal of Tropical Disease*.2012;99-102