



**IJPPR**

INTERNATIONAL JOURNAL OF PHARMACY & PHARMACEUTICAL RESEARCH  
An official Publication of Human Journals

ISSN 2349-7203




Human Journals

Research Article


June 2020 Vol.:18, Issue:3

© All rights are reserved by BAKOU Niangoran François et al.

## The Hydroalcoholic Extract of *Rauvolfia vomitoria* Induces Anxiolytic-Like Effects in Rats



**IJPPR**  
INTERNATIONAL JOURNAL OF PHARMACY & PHARMACEUTICAL RESEARCH  
An official Publication of Human Journals



**BAKOU Niangoran François<sup>\*1</sup>, BA Abdoulaye<sup>2</sup>,  
GUIRO Hamidou<sup>2</sup>, ATAYI E.<sup>3</sup>**

*1 Unit of Animal Physiology, Jean Lorougnon GUEDE  
University, Daloa, (Côte d'Ivoire)*

*2 Laboratory of Neuroscience, UFR Biosciences, Felix  
HOUPHOUET-BOIGNY University, Abidjan, (Côte  
d'Ivoire)*

*3 Neurology Service, Functional Exploration Unit of the  
Nervous System, C.H.U. from Cocody-Abidjan, (Côte  
d'Ivoire)*

**Submission:** 25 May 2020  
**Accepted:** 02 June 2020  
**Published:** 30 June 2020

**Keywords:** *Rauvolfia vomitoria*, leaves, rat, anxiety

### ABSTRACT

**Objective:** The aim of the present work was to evaluate the anxiolytic effect of hydroalcoholic extract of leaves from *Rauvolfia vomitoria* in rat. **Materials and Methods:** The hole-board test, elevated plus-maze paradigm and open field test, were studied to assess the anxiolytic activity of hydroalcoholic extract of leaves from *Rauvolfia vomitoria*. The extract of *Rauvolfia vomitoria* (5, 10, and 25 mg/kg, i.p.) and diazepam (1 mg/kg, i.p.) were administered 30 min before the tests. **Results:** The results showed that extract of R.V (10 and 25 mg/kg, i.p) significantly increased the number and duration of head poking in the hole-board test. In the elevated plus-maze, the extract significantly increased the exploration of the open arm in similar way to that of diazepam. At a dose of 10 and 25 mg/kg i.p. the extract significantly increased both the time spent in and the entries into the open arm by rat. Further, in the open field test, the extract significantly increased rearing, assisted rearing, and number of squares traversed, all of which are demonstrations of exploratory behavior. **Conclusion:** The results of the present study suggest that a hydroalcoholic extract of *Rauvolfia vomitoria* leaves may possess an anxiolytic effect.



HUMAN JOURNALS

[www.ijppr.humanjournals.com](http://www.ijppr.humanjournals.com)

## INTRODUCTION

Anxiety disorders are among the most common psychiatric disorders that affect all age groups of the general population [1]. Approximately 450 million people suffer from a mental or behavioral disorder [2]. It is a feeling of apprehension, uncertainty, and fear characterized by physical symptoms such as palpitations, sweating, and feelings of stress [3]. These disorders are widely treated with benzodiazepine anxiolytic agents. However, the clinical use of benzodiazepines is limited by their side effects such as respiratory depression, motor coordination deficits, memory, cognitive dysfunctions, and dependence liability [4, 5]. Therefore, finding novel therapeutic agents with fewer complications in the treatment of anxiety disorder is of major interest to researchers [6]. Medicinal plant with traditional background of use in neurological diseases could be good candidates to find new anxiolytic agents. *Rauwolfia vomitoria* is a tropical shrub which belongs to the family of Apocynaceae. In Africa, the herbal preparations have been made from various parts of this plant. However, the root, root bark, and bark of stem of this plant have been used extensively, particularly for their aphrodisiac, antipsychotic, abortive, insecticidal, dysenteric, anthelmintic, astringent, cardiotonic, diaphoretic, emetic, expectorant, haemostatic, hypotensive, vulnerary, and febrifugic properties [7]. They are also used in traditional medicine to treat a variety of illnesses such as fever, general weakness, gastrointestinal diseases, liver diseases, psychosis, pain, and cancers [8]. The extract of *R. vomitoria* has been used as traditional medicine for over 2000 years in Africa for the treatment of hypertension and mental disorders and recent studies have confirmed its effectiveness as antipsychotic, antihypertensive, and anti-inflammatory agent and for improving blood chemistry. The anxiolytic property remains to be investigated. The present study was undertaken to assess the possible anxiolytic effects following single administration of hydro-ethanolic extract of *Rauwolfia vomitoria*, Afz in rats. For this purpose, we used the elevated plus-maze, open field and Hole board tests.

## MATERIALS AND METHODS

### Plant material

Fresh leaves of the plant were collected from Daloa, (Cote d'Ivoire) in October 2019. The plant was identified and verified by botanist Professor from Jean Lorougnon GUEDE University of Daloa (Cote d'Ivoire). The collected leaves were dried under a shade during

two weeks and pulverized using the crushing assistance (IKAMAG RCT®). The powder of leaves obtained, constituted our sample to be analyzed.

### **Extract preparation**

A 100 g sample of crushed *Rauwolfia vomitoria* was used for extraction. The sample was soaked overnight in 70% alcohol (30:70) and filtered using Whatman No.1 paper. The process was repeated twice by adding fresh solvent every time. The pooled extract was subjected to flash evaporation followed by lyophilization. The lyophilized sample was further analyzed to determine its anxiolytic property.

### **Animals**

Male Wistar rats aged 8-10 weeks weighing (145 - 250 g) were obtained from the animal house of Jean Lorougnon GUEDE University, Daloa. These animals were housed under standard environmental conditions. The rats were fed with FACI® (Fabrication d'Aliments de Côte d'Ivoire) pellets, groundnuts and dried fish. They had free access to drinking water ad libitum.

### **Drugs and chemicals**

The standard drugs Diazepam was collected from Square Pharmaceuticals Ltd., Cote d'Ivoire. Saline water which was used for dilution purpose was prepared was obtained from Jean Lorougnon GUEDE university of Daloa (Cote d'Ivoire).

### **Behavioral parameters used to test anxiolytic activity**

#### **Open field test**

Locomotor activity and exploratory behavior were assessed in an open field by the method described by Souza [9]. The apparatus consisted of a wooden box ( $60 \times 60 \times 30 \text{ cm}^3$ ) with the floor divided into 16 squares ( $15 \times 15 \text{ cm}^2$ ). The apparatus was illuminated with a 40-W lamp suspended 100 cm above. Twenty-five rats were randomly divided into five groups of five rats each. One hour before test session, rats were treated with graded doses of RV (5, 10, and 25 mg/kg, i.p.) while the control received 10 ml normal saline/kg i.p. 30 min later each rat was placed individually in the center of the apparatus and observed for 5 min to record the

locomotor (number of squares crossed with four paws) and exploratory activities (indicated by frequency of rearing and assisted rearing) [9].

### **Elevated plus-maze test**

The elevated plus maze is an anxiety paradigm based on the rodent's natural aversion to a novel and potentially dangerous environment represented by the open and elevated spaces [10]. The elevated plus maze apparatus is a plus (+) shaped wooden structure, consisting of two open arms ( $40 \times 5 \times 10 \text{ cm}^3$ ) and two enclosed arms ( $40 \times 5 \times 10 \text{ cm}^3$ ) extended from a central platform ( $10 \times 10 \text{ cm}^2$ ). The maze was elevated 50 cm from the room floor. Rats were randomly divided into five groups. The rats that served as control group received 10 ml normal saline/kg body weight i.p, while the treated rats received RV (5, 10, and 25 mg/kg body weight i.p) and diazepam (1 mg/kg body weight i.p.). Thirty minutes after intraperitoneal administration of diazepam, each rat was placed at the center of the maze, facing one of the open arms and allowed to explore the maze freely for a 5-min testing period. The time spent in open and enclosed arms were recorded. The maze was thoroughly cleaned between tests with a tissue paper moistened with 70% ethanol.

### **Hole board test**

The hole board apparatus consisted of a wooden chamber ( $40 \times 40 \times 25 \text{ cm}^3$ ) with 16 holes (each of 3 cm diameter) evenly distributed on the floor. The apparatus was elevated to a height of 25 cm from the ground so that the rat could peep through the holes. The rats were treated with RV (5, 10, and 25 mg/kg body weight i.p), diazepam (1 mg/kg body weight i.p) or distilled water (i.p) 30 min prior to test and kept in the apparatus. The numbers and the duration of head poking were recorded during the 5 min observation period.

### **Statistical analysis**

Results were expressed as mean  $\pm$  S.E.M. The statistical analysis of data was done using the one-way analysis of variance (ANOVA) followed by Dunnett's test. A probability level less than 0.05 was considered statistically significant.

## RESULTS

### Open field test

There was significant ( $*P<0.05$ ) increase in rearing, assisted rearing and number of squares traveled in the *R.V* treated (5 to 25mg/kg i.p.) as well as the standard group, compared to the control group, in the open field test. The numbers of assisted rearing in the groups treated with *R.V* at 5 and 10 mg/kg was comparable with the standard and at 25 mg/kg, the count was significantly ( $*P<0.05$ ) higher than the standard group. The number of squares traveled also significantly ( $*P<0.05$ ) increased when *R.V* was administered at 5,10 and 25 mg/kg, showing anxiolytic activity of the plant extract. (Table 1)

**Table No. 1: Effect of RV on rearing, assisted rearing and squares traveled in open field test (n=5). Values are expressed as mean  $\pm$  SEM**

Treatment	Rearing	Assisted rearing	Number of square traversed
NaCl 10ml	3.5 $\pm$ 1.2	6 $\pm$ 1.6	5 $\pm$ 1.8
DZP 1 mg/Kg	6.1 $\pm$ 2.3*	12 $\pm$ 1.8*	30 $\pm$ 2.6*
RV 5 mg/Kg	6.75 $\pm$ 2.6*	13.66 $\pm$ 2.1*	12 $\pm$ 2.1*
RV 10 mg/Kg	7.33 $\pm$ 5.2*	14.2 $\pm$ 1.9*	14.5 $\pm$ 2.3*
RV 25 mg/Kg	8.5 $\pm$ 6.4*	14.33 $\pm$ 2.3*	15.2 $\pm$ 2.6*

### Elevated plus maze test

In this test, there was significant ( $*P<0.05$ ) increase in the number of entries and the time spent in the open arm *RV* (5 to 25 mg/kg, i.p) and diazepam treated groups, whereas there was a significant ( $*P<0.05$ ) decrease in the time spent in the close arm compared to the control group. Numbers of entries into the open arm in *R.V* treated animals with 25mg/kg dose were higher than the standard drug. (Table 2)

**Table No. 2: Effect of RV on time spent (s) in open arm, time spent (s) in closed arm, entries in open arm and entries in closed arm in elevated plus maze test (n=5). Values are expressed as mean  $\pm$  SEM**

Treatment	Time spent in the open arm (s)	Time spent in the enclosed arm (s)	Entries into open arm	Entries into enclosed arm
NaCl 10 ml	38 $\pm$ 2.6	265 $\pm$ 11.2	4.31 $\pm$ 1.25	12.21 $\pm$ 1.6
DZP 1 mg/Kg	91 $\pm$ 4.16*	56 $\pm$ 2.6*	6 $\pm$ 1.6*	11.8 $\pm$ 1.2
RV 5 mg/Kg	85 $\pm$ 3.6*	245 $\pm$ 10.6	5.17 $\pm$ 2.01	11 $\pm$ 1.13
RV 10 mg/Kg	87 $\pm$ 3.8*	210 $\pm$ 11.4	7.31 $\pm$ 2.03*	9 $\pm$ 2.01
RV 25 mg/Kg	89 $\pm$ 4.01*	77 $\pm$ 3.06*	8 $\pm$ 2.6*	7.6 $\pm$ 2.2*

### Hole board test

There was significant (\*\*P<0.01) and dose-dependent increase in the number and duration of head poking after administration of RV (10 and 25 mg/kg, i.p) compared to the control group and the results were comparable with the standard drug diazepam. (Table 3).

**Table No. 3: Effect of RV on head poking and duration (s) of head pokes in hole board test (n=5). Values are expressed as mean  $\pm$  SEM**

Treatment	Dose route	Duration of head poking (s)	Number of head poking
NaCl	10 ml	30 $\pm$ 1.06	35 $\pm$ 1.1
DZP	1 mg/Kg	56 $\pm$ 1.8	63 $\pm$ 2.6
RV	5 mg/Kg	36 $\pm$ 1.12	38 $\pm$ 1.6
RV	10 mg/Kg	60 $\pm$ 2.5**	55 $\pm$ 1.6**
RV	25 mg/Kg	67 $\pm$ 2.7**	65 $\pm$ 2.4**

### DISCUSSION

The open-field apparatus provides information on anxiety-related behaviour characterized by natural aversion of rodents to an open brightly lit area [11]. Animals are thus afraid of the centre and spend more time in the protective corners and in freezing state. Anxiolytics increase total locomotive activity resulting in a reduction of time spent in corners, increased time spent in the center and a decreased time spent in freezing state. The extract of *R. V* at 5, 10 and 25 mg/kg body weight increased total locomotive activity and increased rearing of

treated rats in our study. The elevated plus maze (EPM) test represents one of the most widely used animal models for screening anxiolytics [10]. This test is able to reproduce anxiolytic or anxiogenic effects in rodents such that anxiolytics produce increase the time spent in the open arm of the elevated plus maze, while anxiogenic substances produce the opposite effect [10, 12]. The indices of anxiety (number of open-arm entries, and time spent in the open arm) are sensitive to agents and are thought to act via the GABAA receptor complex, justifying the use of diazepam (DZP) as a positive control in this study. Diazepam, a benzodiazepine binds to GABAA receptors to increase the frequency of chloride channel openings resulting in hyperpolarization. It increased the frequency of open-arm entries and the time spent in the open arms [13], confirming its anxiolytic effects. In our study, we observed that *R.V* (5, 10 and 20 mg/kg) induced significant increases in both the number of entries and time spent in the open arms. The number of entries and the time spent in the closed arms were reduced in the extract-treated group as compared to the control group. Hole-board test indicated that the head-dipping behavior was sensitive to changes in the emotional state of the animal, and suggested that the expression of an anxiolytic state may be reflected by an increase in head-dipping behavior [14]. In our study, *R.V* (5,10 and 20mg/kg) significantly increased the numbers and duration of head poking compared to the control group. These results confirm the anxiolytic effects of *Rauvolfia vomitoria*. They are to be compared with the work of [15] Nsour, who in a similar study showed the anxiolytic effect of *Rauvolfia serpentina*.; [16] from Aidee, which highlighted the anxiolytic effects of the ethanolic extracts of *Argemone mexicana* ; [17] from Carla, which demonstrated anxiolytic properties of aqueous extracts of *Salvia miltiorrhiza* in rats; [18] Charles and [19] Carnevale which showed anxiolytic properties of extracts of *Maerua angolensis* in mice and *Griffonia simplicifolia* in rat. The anxiolytic effect of the hydroethanolic leaves from *Rauvolfia vomitoria* could be due to the presence of alkaloids among the compounds of *Rauvolfia vomitoria*. Indeed, [16] Aidee demonstrated that the alkaloids isolated from *Argemone mexicana* extracts increased the percentage of time spent in the open arms of rat EPM, in the same way as diazepam and *Argemone mexicana* extracts.

## CONCLUSION

In conclusion, our results showed that the hydroethanolic leaves of *Rauvolfia vomitoria*, have anxiolytic-like effects in rat. This property possibly due to the presence of different phytoconstituents like alkaloids, steroids and present therein. However, the exact mechanism



(s) related to the active ingredient (s) in *Rauvolfia vomitoria* extract should be elucidated in future studies.

## REFERENCES

- 1 Kessler RC, Angermeyer M, Anthony JC, Lifetime prevalence and age-of-onset distributions of mental disorders in the World Health Organization's World Mental Health Survey initiative. *World Psychiatry* 2007, 6: 168–76.
- 2 World Health Report. Mental health: New understanding new hope. WHO, Geneva. 2001.
- 3 Silvaraman D, Muralidharan P, Habibur R. Evaluation of the anxiolytic effect of methanolic leaf extract of *Ficus hispida* Linn. in corticosterone induced anxiety in young adult rats. *Pharmacologia*, 2012, 3: 467-471.
- 4 Venault P, Chapouthier G, de Carvalho LP, Benzodiazepine impairs and beta-carboline enhances performance in learning and memory tasks. *Nature* 1986, 321: 864-6.
- 5 Greenblatt DJ and Shader RI. 1978 Dependence, tolerance, and addiction to benzodiazepines: clinical and pharmacokinetic considerations. *Drug Metab. Rev.* 1978, 8:13-28.
- 6 Kent JM, Mathew SJ and Gorman JM. Molecular targets in the treatment of anxiety. *Biol. Psychiatry* 2002, 52: 1008-30.
- 7 Eluwa M., Idumesaro N., Ekong M., Akpantah A., Ekanem T. Effect of aqueous extract of *Rauvolfia vomitoria* root bark on the cytoarchitecture of the cerebellum and neurobehaviour of adult male wistar rats. *The Internet Journal of Alternative Medicine*; 2008, 6 (2):1–7.
- 8 Yu J., Chen Q. Antitumor activities of *Rauvolfia vomitoria* extract and potentiation of gemcitabine effects against pancreatic cancer. *Integrative Cancer Therapies*; 2014,13(3):217–225.
- 9 Souza G, Christina A, Cesar AB, WN. Diphenyl diselenide improves scopolamine-induced memory impairment in mice. *Behav Pharmac*, 2010, 21: 556-562.
- 10 Lister RG, 1987. The use of a plus-maze to measure anxiety in the mouse. *Psychopharmacol (Berl)*, 1987, 92: 180-185.
- 11 Choleris E, Thomas AW, Kavaliers, A detailed ethological analysis of the mouse open field test: Effects of diazepam, chlordiazepoxide and an extremely low frequency pulsed magnetic field. *Neurosci Behav Rev*, 2001, 25: 235-260.
- 12 Pellow S, File SE. Anxiolytic and anxiogenic drug effects on exploratory activity in an elevated plus maze: a novel test of anxiety in rats. *Pharmacol Biochem Behav*, 1986, 24: 525-529.
- 13 Crawley JN. Behavioral phenotyping of transgenic and knockout mice: Experimental design and evaluation of general health, sensory functions, motor abilities and specific behavioural tests. *Brain Res*, 1999, 835: 18-26.
- 14 Chandana C.B., Archana T., Shameem A.B. Anxiolytic activity of methanol leaf extract of *Achyranthes aspera* Linn in mice using experimental models of anxiety. *Indian J Pharmacol.*, 2012; 44(1): 63–67.
- 15 Nsour W. M., Lau C. B., Wong I. C. Review on phytotherapy in epilepsy Seizure; 2001, 9: 96–107.
- 16 Aideé I. A, Omar D. M., Miguel Á. D. Anxiolytic-like effect of ethanolic extract of *Argemone mexicana* and its alkaloids in Wistar rats 2016, 6:476-488.
- 17 Carla L., Giancarlo C., Gian L. G. anxiolytic effect of an extract of *Salvia miltiorrhiza* roots in rats 2018, 81: 390-397
- 18 Charles K. B., Robert P. B., Donatus W. A, Anxiolytic and Antidepressant Effects of *Maerua angolensis* DC. Stem Bark Extract in Mice. 2018, 5:1-17
- 19 Carnevale G., Di Viesti V., Zavatti M. Anxiolytic-like effect of *Griffonia simplicifolia* Baill. Seed extract in rats. *Phytomedicine*, 2011,18(10):848-851