



# IJPPR

INTERNATIONAL JOURNAL OF PHARMACY & PHARMACEUTICAL RESEARCH  
An official Publication of Human Journals

ISSN 2349-7203



Human Journals

Research Article

February 2021 Vol.:20, Issue:3

© All rights are reserved by Kapil Jangir et al.

## Evaluation of Anti-Pyretic Activity of *Woodfordia fruticosa* Using Brewer's Yeast Induced Pyrexia in Albino Rats

	
<b>Kapil Jangir*, Nitin Jumnani</b>	
<i>Alwar Pharmacy College, Alwar - 301030, Rajasthan, India</i>	
<b>Submitted:</b>	03 January 2021
<b>Revised:</b>	23 January 2021
<b>Accepted:</b>	12 February 2021



[www.ijppr.humanjournals.com](http://www.ijppr.humanjournals.com)

**Keywords:** Anti-pyretic activity, *Woodfordia fruticosa* (Dhataki), Paracetamol, Yeast, albino rats.

### ABSTRACT

**Background:** Many herbal plants are having antipyretic effects. *W. fruticosa* traditionally used to treat fever pain and inflammation. **Aim:** The present study investigates the antipyretic activity of the brownish extracts of *Woodfordia fruticosa* flower on brewer's yeast induced fever in experimental rats. **Methods:** 30 albino rats weighing 150-180g were used for study. Brewer's-yeast-induced pyrexia was used to investigate the antipyretic effects. 10ml/Kg (subcutaneous) of 0.5(w/v) suspension of brewer's yeast in Carboxymethylcellulose was injected to induce fever in all the experimental animals. After 18hrs, the rectal temperature was taken and the animals were administered *Woodfordia fruticosa* (100mg/kg, 200mg/kg, and 400mg/kg) and paracetamol (standard group, 30mg/kg) orally. **Result:** The body temperature of the rats was measured rectally over a period of 3hours. *W. fruticosa* (200mg/kg and 400mg/kg) significantly reduced yeast-induced pyrexia when compared with the standard drug. Thus, this experiment shows that the antipyretic effect of *W. fruticosa* is dose-dependent and the effect is as a result of the flavonoid component of the extract. **Conclusion:** These data therefore suggest that extract of *W. fruticosa* possesses significant antipyretic activity and its mechanism could be by inhibition of release inflammatory mediators and prostaglandins.

## INTRODUCTION:

Over the centuries, plants have been used for their medicinal values. However, many plants have yet to be studied and relatively few scientific researches have been carried out.<sup>1, 2</sup> Therefore, assessment of medicinal plants is of special interest because of rich heritage and the continuous use among large portion of people.<sup>1-3</sup>

Fever, is known as pyrexia and may occur due to infection, inflammation, or any tissue damage and disease states. It is defined as the elevation of core body temperature above normal; in normal adults, the average oral temperature is 37°C (98.6°F) (1). Fever is a medical sign rather than a disease and it may be caused as a secondary impact of infection, malignancy or other diseased states. Fever occurs when the body's thermostat (located in the hypothalamus-anterior pituitary) resets at a higher temperature, primarily in response to an infection. From the recent scientific discovery, most of the antipyretic drugs have been developed to reduce elevated body temperature, of which many acts by the mechanism of inhibition of the COX-2 expression to reduce PGE2 biosynthesis.<sup>4-7</sup>

It is currently accepted that prostaglandin E2 (PGE2) is the final fever mediator in the brain, specifically in the pre-optic area of the anterior hypothalamus. Many drugs on chronic usage can able to result in several side effects including gastrointestinal, renal, hepatic, central nervous system and dermatological effects. A rich heritage of knowledge on preventive and curative medicines was available in ancient scholastic work.<sup>4, 5, 7</sup> The beneficial medicinal effects of plant materials typically result from the combinations of secondary products present in the plant. Many of the plants were found to show the antipyretic activity hence the use of traditional plants as relieving agents can be advantageous in many aspects such as easy availability, economical with no or limited side effects.<sup>8</sup>

## MATERIALS AND METHODS:

*Woodfordia fruticosa*, also called **Dhai**, Dhataki, Fire Flame Bush is a species of plant from the Lythraceae family. In Ayurveda this herb is used to treat many diseases. Its flower is analgesic antipyretic and astringent due to pitta dosha. It is widely cultivated in tropical regions, usually between 0 and 300 m altitude, and grows well in brushwood, bush margins, grassy fields, and secondary forest, seasonally flooded grassy fields and sandy coasts and along roadsides and riverbanks, although the species cannot tolerate excessive water or heat.<sup>9-</sup>

17

**Plant Taxonomy:** <sup>18-21</sup>

**Table No. 1: Scientific classification of *Woodfordia fruticosa***

<b>Synonyms</b>	Dhai, Dhati, Dhataki, Dawai, Dhavi
<b>Kingdom</b>	Plantae
<b>Subkingdom:</b>	Tracheobionta
<b>Division:</b>	Magnoliophyta
<b>Class:</b>	Magnoliopsida
<b>Subclass:</b>	Rosidae
<b>Order:</b>	Myrtales
<b>Family:</b>	Lythraceae
<b>Genus:</b>	<i>Woodfordia</i>
<b>Species:</b>	<i>fruticosa</i>

**Drugs and chemicals:**

Paracetamol was purchased from CDFCL. The solvents and other chemicals of analytical grade were used and obtained from the institute's central store. Brewer's yeast was purchased from local market.

**Collection & extraction of Flowers:**

The fresh flowers of *W. fruticosa* were collected from the local areas of Alwar district and are air dried under shade for two days and then coarsely powdered.

**Defattation of powdered *Woodfordia fruticosa* flower**

200 g coarse flower powdered were defatted with 400 ml petroleum ether (40-60°C) using soxhlet apparatus. Extraction was continued until a drop of solvent from siphon tube, when evaporated on filter paper, did not leave a greasy spot (approximately 10-12 cycles). After the defattation, mark was taken out from extractor and spreaded as a bed on a clean paper and dried till evaporation of petroleum ether.<sup>22-24</sup>

**Ethanollic extraction of *W. fruticosa*:**

The dried mark obtained after defattation was packed in soxhlet apparatus and extracted with 400 ml ethyl alcohol in soxhlet apparatus. Extraction was continued until a drop of solvent from the siphon tube, when taken on TLC plate and sprayed with concentrated sulphuric acid, does not give a black spot. Brownish extract thus obtained, was collected and solvent was evaporated under reduced pressure. The percentage yield of ethanolic extract was found to be 10.12 % which is comparable with the reported standard extractive value.<sup>25-27</sup>

**Animals**

30 albino rats weighing between 150-180g of either sex were used for this experimental study. They were divided into five groups (n=6). To the rats laboratory, diet and water *ad libitum* was provided and were kept in polypropylene cages. Which are kept in a constant humidity (55%), temperature at (22± 2<sup>0</sup>C), and exposed to dark and light { 12hr} every day the bedding materials of the cages were changed. The experimental protocol was approved by IAEC, with proposal no APC/CPCSEA/2020/1709, for using animals in this experiment. Animals were fasted overnight with free access to water before each experiment.<sup>24, 28</sup>

**Table No. 2: Grouping and dose of drug for anti-pyretic activity**

Group	Drug and Doses
<b>Brewer’s yeast-induced pyrexia</b>	
Group I	Normal Control (Received 0.5% CMC)
Group II	<i>W. fruticosa</i> 100mg (i.p.)
Group III	<i>W. fruticosa</i> 200mg (i.p.)
Group IV	<i>W. fruticosa</i> 400mg (i.p.)
Group V	<i>Paracetamol</i> 30 mg/kg (oral)

**Induction of Pyrexia:**

Fever was induced by injecting 10ml/Kg (subcutaneous) of 0.5(w/v) suspension of brewer’s yeast in Carboxy methylcellulose below the nape of the neck. The temperature was measured after 18hours using rectal thermometer.<sup>29, 30</sup>

**Drug administration:**

After 18 h of yeast injection 0.5% CMC was administered orally to the control Group I. The extracts of *W. fruticosa* were administered at doses of (100, 200 and 400 mg/kg b.w. p.o) to groups II, III, IV animals respectively. Group V animals received the standard drug Paracetamol (30 mg/kg b.w. p.o) and Rats were restrained for recording rectal temperature by at the 18 h, immediately before extracts, normal saline or paracetamol administration, and again at one-hour intervals up to the 21 h after yeast injection.

**Statistical Analysis:**

Results are expressed as mean ± SEM. Statistical analysis of data was performed using ANOVA to study differences among the means.

**RESULTS AND DISCUSSION:**

Fever is known to be caused by several endogenous pyrogens such as interleukin-1β, interleukin-6, interleukin-8, tumor necrosis factor-α, macrophageprotein-1 and prostaglandins. Prostaglandin synthesis may be activated by tumor necrosis factor-α and phospholipase A2. Brewer’s yeast induces both TNF-α and prostaglandin synthesis.

**Table No. 3: Antipyretic Effect of Ethanolic extract of *Woodfordia fruticosa* on yeast induced pyrexia in rats.**


Treatment	Dose mg/kg	Rectal temp. °C before and after treatment						
		Normal	18h	30 min after treatment	60 min after treatment	90 min after treatment	120 min after treatment	180 min after treatment
Control	-	37.7±0.18	38.7±0.34	38.6±0.21	38.5±0.04	38.5±0.24	38.5±0.12	38.5±0.16
Paracetamol	30	37.6±0.12	38.6±0.24	38.2±0.13	37.8±0.09	37.7±0.04	37.6±0.26	37.62±0.06
EEWF	100	37.7±0.17	38.6±0.32	38.2±0.04	38.12±0.05	37.98±0.16	37.90±0.07	37.82±0.05
EEWF	200	37.6±0.21	38.6±0.18	38.12±0.05	37.9±0.05	37.82±0.04	37.7±0.04	37.7±0.06
EEWF	400	37.6±0.08	38.6±0.15	37.9±0.06**	37.8±0.03**	37.8±0.06**	37.68±0.08**	37.67±0.07

Values are expressed as mean ± SEM (n=6). \*\*P<0.01 Statistically.

The subcutaneous injection of yeast suspension (Nape of neck) markedly elevated the rectal temperature after 18 h of administration. Treatment with the WF at doses 200 and 400 mg/kg significantly ( $p < 0.01$ ) decreased the rectal temperature of the rats in dose dependent manner up to 3 h after pyrexia when compared with control (Table 3). Effect of 100mg/kg extract was negligible. The antipyretic effect was sustained throughout the remaining period of the experiment in a manner similar to the reference drug Paracetamol 30mg/kg, b.w. p.o.

The extracts are likely to reduce fever by reducing brain concentration of prostaglandin E2 especially in the hypothalamus through its action on COX-2 or by enhancement of the production of the body's own antipyretic substances.

The efficacy of the antipyretic effect of dhataki extract was observed to have increased with increased concentration (dose-dependent manner). This can be said to be due to the increased concentration of the component of the extract exhibiting antipyretic effects.



**Figure No. 1: Graphical Representation of Anti-pyretic Effect of Ethanolic extract of *W. fruticosa* on yeast induced pyrexia in rats**

**CONCLUSION:**

Antipyretic activity of bioflavonoid from *Woodfordia fruticosa* plant extract was confirmed by the present experimental studies. *W. fruticosa* extract with two different doses of 200mg/kg and 400mg/kg showed the reduced temperature levels in the wistar albino

experimental rats when compared to the temperatures of rats of control group. It is through the inhibition of prostaglandin (PGE<sub>2</sub>). From these results, it can be concluded that the *W. fruticosa* plant extract at the dose level of 200mg/kg has shown a marked decrease in temperature to normal. Hence possess antipyretic activity which was comparable with the paracetamol effect. Plant test extract of 100mg/kg was not so efficient as the dose level of 200mg/kg.

### CONFLICT OF INTERESTS:

The authors declare that there is no conflict of interests regarding the publication of this paper.

### REFERENCES:

1. Benzie FF, Galor SW, "Herbal Medicine-Biomolecular and Clinical Aspects", 2nd edition, 2011, 114-123.
2. Yuan H, Piao G, "The Traditional Medicine and Modern Medicine from Natural Products", *Molecules*, 2016, 21, 559, 1-18.
3. Pathare YS, Wagh VD, "Herbal Medicines and Nutritional supplements used in the treatment of Glaucoma: A Review", *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 2012, 3(1), 331-337.
4. Ogoina D, "Fever, fever patterns and diseases called 'fever' - A review", *ELSEVIER-Journal of Infection and Public Health*, August 2011, 4(3), 108-124.
5. Walter EJ, Jumma SH, Carraretto M, "The pathophysiological basis and consequences of fever", *Critical Care*, 2016, 20:200, 1-10.
6. Nakamura K, "Central circuitries for body temperature regulation and fever", *Am J Physiol Regul Integr Comp Physiology*, 2011, R1207-R1218.
7. Jeffrey RA, Baker MD, "Management of fever in infants and children", *Emergency Medicine Clinics of North America*, Mar 2002, 20(1):49-67
8. Garg V, Dutt R, "Antipyretic Plants: An Updated Review", *Current Bioactive Compounds*, 2020, 16 (1), 4-12.
9. Vanmala VB, Archana G. Mhase *et al*, "Preliminary pharmacognostic and phytochemical standardization of Dhataki [*Woodfordia fruticosa* (L.) Kurz.] leaves", *Ayurved*, 2014, 35(3): 309-315.
10. Khare CP, *Indian Medicinal Plants*, published by Springer-Verlag Heidelberg, 2004; 328-329.
11. Kirtikar KR, Basu BD, *Indian Medicinal Plants*, Published by International Book Distributors Dehradun India, Vol-I, 2nd edition; 1998, 712.
12. Atal CK, Bhatia AK, Singh RP, "Role of *Woodfordia fruticosa* Kurz in the preparation of asavas and arishtas", *Journal of Research in Ayurveda and Siddha*, 1982, 3: 193-199.
13. Chauhan JS, Srivastava SK, Srivastava SD (1979a). Phytochemical investigation of the flowers of *Woodfordia fruticosa*. *Planta Medica* 36:183-184.
14. Das PK, Goswami S, Chinniah A, Panda N, Banerjee S, Sahu NP, Achari B (2007). *Woodfordia fruticosa*: Traditional uses and recent findings. *Journal of Ethnopharmacology* 110: 189-199.
15. Galor SW, Benzie IF, "Herbal Medicine: A Growing Field With A Long Tradition", *Herbal Medicine: Biomolecular and Clinical Aspects*, 2<sup>nd</sup> Edition 2011, 1-5.
16. Janghel V, Chandel SS, Venkatachalam T, "Preliminary phytochemical and physicochemical investigation of *Woodfordia fruticosa* (linn) kurz root", 2019, 12 (4), 840-845.
17. Syed YH, Khan M, Bhuvaneshwari J, "Phytochemical investigation and standardization of extracts of flowers of *Woodfordia fruticosa*; a preliminary study", *Journal of Pharm. Bio Science*, 2013, 4, 134-140.
18. Kumar D, Sharma M, Sorout A, "Woodfordia fruticosa Kurz.: A Review on its Botany, Chemistry and Biological activities", *Journal of Pharmacognosy and Phytochemistry* 2016; 5(3): 293-298.



19. Srivastava AK, Siddharth SA, Nagar H, "Phytopharmacological evaluation of aerial parts of *Woodfordia fruticosa* (L.) Kurz in Cough Variant Asthma" Pharmacognosy Journal, Sep-Oct, 2015, 7(5), 296-299.
20. Verma N, Amresh G, Sah PK, "Anti-inflammatory and antinociceptive activity of hydroethanolic extract of *Woodfordia fruticosa* Kurz flowers", Pelagia Research Library, Der Pharmacia Sinica, 2012, 3(2):289-294.
21. Ghante MH, Bhusari KP, Duragkar NJ, "Pharmacological evaluation for anti-asthmatic and anti-inflammatory potential of *Woodfordia fruticosa* flower extracts", Pharmaceutical Biology, 2014, 1-10.
22. Hemraj, Upmanyu N, Gupta A, "Pharmacological Activities of *Stephania glabra*, *Woodfordia fruticosa* and *Cissempeles pareira*-A Review", International journal of pharmacy and pharmaceutical sciences, 2012, 4(3), 16-23.
23. Mujalde VS, "Role of Dhatakiphuspa (*Woodfordia fruticosa* Kurz) in pravahika", Global Journal for Research Analysis, July 2019, 8(7), 155-156.
24. Uday M, Kishor D, Ajay R, "A Pharmacognostic and pharmacological overview on *Woodfordia fruticosa* Kurz.", Scholars Academic Journal of Pharmacy, 2014; 3(5): 418-422.
25. Shubhashre MN, Shantha TR, Ramarao V, "A Review on Therapeutic Uses of Flowers as Depicted in Classical Texts of Ayurveda and Siddha", J. Res. Educ. Indian Med, March 2015, Vol -21, 1-5.
26. Sharma S, Sharma N, "Pharmacognostical Evaluation of Dhataki (*Woodfordia fruticosa* Kurtz.)", International Journal of Recent Scientific Research, 2019, 10(9), 35139-35143.
27. Das PK, Goswami S, "*Woodfordia fruticosa*: Traditional uses and recent findings", Journal of Ethnopharmacology ELSEVIER, 2007, 189-199.
28. Parekh J, Chanda S, "In Vitro Antibacterial activity of the crude methanol extract of *Woodfordia fruticosa* kurz. Flower (lythraceae)", Brazilian Journal of Microbiology 2007, 38, 204-207.
29. Tomazetti J, Avila DS, Ferreira A *et al.*, "Baker yeast-induced fever in young rats: Characterization and validation of an animal model for antipyretics screening", ELSEVIER Journal of Neuroscience Methods, 2005, 147; 29-35.
30. Kumar SA, Venkatarathanamma V, "Antipyretic Activity of Annona Plants Leaves on Brewer's Yeast Induced Febrile Rats", Asian J Pharm Clin Res, 2015, 8(3), 210-212.

