



IJPPR

INTERNATIONAL JOURNAL OF PHARMACY & PHARMACEUTICAL RESEARCH
An official Publication of Human Journals

ISSN 2349-7203




Human Journals

Review Article


May 2023 Vol.:27, Issue:2

© All rights are reserved by Pankaj Sharma et al.

Synthesis, Structure Activity Relationship and Pharmaceutical Applications of Pyrimidine Analogues: A Review



IJPPR
INTERNATIONAL JOURNAL OF PHARMACY & PHARMACEUTICAL RESEARCH
An official Publication of Human Journals



**Pankaj Sharma^{1*}, Abhishek Soni², Muskan Sharma³,
Madhurima Thakur⁴, Sheetal Rani⁵**

1Research Scholar, Department of Pharmaceutical Chemistry, School of Pharmacy, Abhilashi University, Chail Chowk (H.P.) India.

2Associate Professor, Department of Pharmaceutics, Abhilashi College of Pharmacy. Nerchowk, Mandi (H.P.). India.

3Research Scholar, Department of Pharmaceutical Chemistry, School of Pharmacy, Abhilashi University, Chail Chowk (H.P.) India.

4Research Scholar, Department of Pharmaceutics, School of Pharmacy, Abhilashi University, Chail Chowk (H.P.) India.

5Research Scholar, University Institute of Pharmaceutical Science, Chandigarh University, Gharuan, Mohali, (Punjab). India.

Submitted: 27 April 2023
Accepted: 02 May 2023
Published: 30 May 2023

Keywords: Pyrimidine, analogues, chemical structure, synthesis, and biological applications

ABSTRACT

Pyrimidine and its analogues are important heterocyclic compounds that have various applications in chemistry and pharmacy. These compounds are composed of a six-membered ring containing four carbon atoms and two nitrogen atoms, which are arranged in a planar fashion. Pyrimidine is a fundamental building block of nucleic acids such as DNA and RNA and plays a critical role in cellular metabolism. Pyrimidine analogues have attracted significant attention from chemists and pharmacologists due to their diverse biological activities, such as antiviral, antibacterial, anticancer, and anti-inflammatory properties. In this article, we provide an overview of the chemical structure, synthesis, and biological applications of pyrimidine and its analogues. We also discuss various classical and modern methods for the synthesis of pyrimidine analogues, including transition-metal-catalyzed reactions, click chemistry, and microwave-assisted reactions. The diverse range of biological activities exhibited by pyrimidine analogues has led to the development of numerous pyrimidine-based drugs, including anticancer agents such as 5-fluorouracil and capecitabine, and antiviral drugs such as acyclovir and ribavirin. The synthesis and biological applications of pyrimidine and its analogues are of great interest to chemists and pharmacologists, and they continue to be an active area of research.




HUMAN JOURNALS

www.ijppr.humanjournals.com

1. INTRODUCTION:

Pyrimidine and its analogues are an essential class of heterocyclic compounds that have found numerous applications in chemistry and pharmacy. These molecules are composed of a six-membered ring containing four carbon atoms and two nitrogen atoms, which are arranged in a planar fashion (Obaid RJ, *et. al.*, 2022). Pyrimidine is a fundamental building block of nucleic acids such as DNA and RNA, and it plays a critical role in cellular metabolism. Pyrimidine derivatives and analogues have attracted significant attention from chemists and pharmacologists due to their diverse biological activities, such as antiviral, antibacterial, anticancer, and anti-inflammatory properties (Mukherjee AK, *et. al.*, 2001). This has led to the development of numerous pyrimidine-based drugs, including anticancer agents such as 5-fluorouracil and capecitabine, and antiviral drugs such as acyclovir and ribavirin (Jain KS, *et. al.*, 2016).

Pyrimidine is a six-membered aromatic ring containing two nitrogen atoms at positions 1 and 3 and four carbon atoms at positions 2, 4, 5, and 6. The chemical formula of pyrimidine is $C_4H_4N_2$, and its molecular weight is 80.09 g/mol. The ring is planar and exhibits a resonance structure in which the double bonds between positions 2 and 4 and between positions 4 and 6 can move between the two nitrogen atoms as shown in Fig. 1. Pyrimidine can be synthesized using various methods, including the Biginelli reaction, the Borsche-Drechsel cyclization, and the Debus-Radziszewski reaction. The Biginelli reaction involves the condensation of an aldehyde or ketone, a urea, and an acid catalyst in the presence of an alcohol solvent to yield a dihydropyrimidine (Pels Rijcken WR, *et. al.*, 1990). The dihydropyrimidine is then oxidized using mild oxidants such as hydrogen peroxide or potassium permanganate to yield the desired pyrimidine. The Borsche-Drechsel cyclization involves the cyclization of an ethyl 2-amino-4, 6-dimethoxypyrimidine-5-carboxylate using phosphorous oxychloride or thionyl chloride to yield pyrimidine-5-carboxylic acid (Choy N, *et. al.*, 2000). The Debus-Radziszewski reaction involves the condensation of an aldehyde, an amine, and a β -keto ester in the presence of an acid catalyst to yield a dihydropyrimidine, which can be oxidized to yield the desired pyrimidine (Stiernet P, *et. al.*, 2019).



Pyrimidine

Fig. 1: Chemical Structure of Pyrimidine

Pyrimidine analogues are compounds with a chemical structure similar to pyrimidine but with different substituents attached to the ring. These compounds can be classified into two categories: pyrimidine nucleosides and pyrimidine derivatives. Pyrimidine nucleosides are compounds made up of a pyrimidine base linked to a sugar molecule (Chiacchio MA, *et. al.*, 2019). These compounds are the building blocks of DNA and RNA and play a critical role in cellular metabolism. Pyrimidine nucleosides include cytidine, uridine, and thymidine, which are essential for the synthesis of DNA and RNA. Pyrimidine derivatives are compounds that contain a pyrimidine ring with one or more substituents attached to the ring (Lagoja IM. *et. al.*, 2005). These substituents can be either alkyl, aryl, or hetero-aryl groups and can significantly alter the biological activity of the compound. Pyrimidine derivatives include a wide range of compounds, such as anticancer agents, antiviral agents, and antihypertensive agents (Keri RS, *et. al.*, 2015).

Pyrimidine is an aromatic heterocyclic compound similar to pyridine. It is one of the three diazines (unsaturated six-membered rings containing two nitrogen atoms) that has two nitrogen atoms at positions 1 and 3 in the ring (Selvam TP, *et. al.*, 2015). Heterocyclic compounds carrying pyrimidine rings are of enormous importance because they represent a vital family of natural and synthetic products, several of which display valuable clinical applications and bioactivities (ur Rashid H, *et. al.*, 2021).

Substituted pyrimidines and purines are extensively found in living things and are among the leading compounds investigated by chemists. Pyrimidines represent the most abundant members of the diazine class, with thymine, uracil, and cytosine being key components of deoxyribonucleic acid and ribonucleic acid (Grazia Martina M, *et. al.*, 2023). Moreover, the pyrimidine moiety occurs in several natural products, for instance, vitamin B1 (thiamine), and various synthetic products, such as barbituric acid and veronal, which are used as soporific drugs as shown in Fig. 2 and Fig. 3.




Fig. 2: Chemical structures of pyrimidine, thymine, uracil, and cytosine




Fig. 3: Chemical structures of thiamine, barbituric acid, and veronal

2. Synthesis of Pyrimidine Analogues:

Pyrimidine analogues can be synthesized using classical or modern methods. Classical methods include the Biginelli, Gewald, Knorr, and Debus-Radziszewski reactions, which involve the condensation of aldehydes, ketones, amines, and carboxylic acids (Gangjee A, *et. al.*, 2004). Modern methods, such as transition-metal-catalysed reactions, click chemistry, and microwave-assisted reactions, are more efficient. Transition-metal-catalysed reactions use transition metal catalysts to form C-C or C-N bonds (Polshettiwar V, *et. al.*, 2008). Click chemistry is a rapid and efficient tool for synthesizing complex molecules, while microwave-assisted reactions utilize microwave radiation to accelerate chemical reactions, resulting in faster reaction times, higher yields, and reduced waste (Pałasz A, *et. al.*, 2015).

The choice of synthetic method depends on the properties of the desired pyrimidine analogue and the goals of the synthesis (Sasada T, *et. al.*, 2009). Some methods of synthesizing pyrimidine analogues are described below:

- **Synthesis via ZnCl₂-catalyzed three-component coupling reaction:** This approach involves a three-component coupling reaction comprising substituted enamines, triethyl orthoformate, and ammonium acetate under ZnCl₂ catalysis, resulting in the synthesis of numerous 4,5-disubstituted pyrimidine analogues in a single step (Foley DJ, *et. al.*, 2020).




Fig. 4: Synthesis via ZnCl₂-catalyzed three-component coupling reaction.

- **Synthesis via K₂S₂O₈-facilitated oxidative annulation reaction:** This approach involves a K₂S₂O₈-facilitated oxidative annulation reaction with formamide as a route towards pyrimidines. Activation of Acetophenone–formamide conjugates resulted in the formation of 4-arylpyrimidines (Jadhav SD, *et. al.*, 2017).



Fig. 5: Synthesis via K₂S₂O₈-facilitated oxidative annulation reaction.

- **Synthesis via Cyclization of ketones with nitriles:** This method involves the synthesis of distinctly substituted pyrimidines through a simple and cost-effective procedure that involves the cyclization of ketones with nitriles under Cu-catalysis in the presence of a base (Su L, *et. al.*, 2018).




Fig. 6: Synthesis via Cyclization of ketones with nitriles

- **Synthesis via reactions of carbonyl compounds with amidine:** This approach involves the regioselective reaction of carbonyl compounds (esters, aldehydes, and ketones) with amidines in the presence of (2, 2, 6, 6-tetramethylpiperidin-1-yl) oxyl (TEMPO) and an in situ prepared recyclable iron (II)-complex. The reactions progress via a TEMPO complexation/ enamine addition/transient a-occupation/b-TEMPO elimination/cyclization order, resulting in the synthesis of numerous pyrimidine analogs (Chu XQ, *et. al.*, 2017).




Fig. 7: Synthesis via reactions of carbonyl compounds with amidine

- Synthesis via base-facilitated intermolecular oxidative C–N bond fabrication:** This method involves a base-facilitated intermolecular oxidative C–N bond fabrication of allylic $\text{C}(\text{sp}^3)\text{--H}$ and vinylic $\text{C}(\text{sp}^2)\text{--H}$ of allylic compounds with amidines to allow easy synthesis of pyrimidines. This approach results in the synthesis of polysubstituted pyrimidines under oxygen as a solitary oxidant, ensuring the supply of protective group-free nitrogen, high efficiency, worthy functional group leniency, and environmental sustainability (Guo W, *et. al.*, 2016).



Fig. 8: Synthesis via base-facilitated intermolecular oxidative C–N bond fabrication

- Synthesis via 4-HO-TEMPO-mediated [3 + 3] annulation of amidines with saturated ketones under Cu-catalysis:** An effective and smooth synthesis of functionally vital pyrimidines has been reported by 4-HO-TEMPO-facilitated [3 + 3] annulation of commercial-grade amidines with saturated ketones under Cu-catalysis. This method provides a new protocol for the synthesis of pyrimidine derivatives via a cascade reaction of oxidative dehydrogenation/annulation/oxidative aromatization using direct beta- $\text{C}(\text{sp}^3)\text{--H}$ functionalization of saturated ketones succeeded by annulation with amidines (Zhan JL., *et. al.*, 2016).




Fig. 9: Synthesis via 4-HO-TEMPO-mediated [3 + 3] annulation of amidines with saturated ketones under Cu-catalysis

- **Synthesis via Viable multicomponent synthesis:** Pyrimidines are also obtained from amidines and up to three (dissimilar) alcohols under iridium-catalysis through a regio-selective, multicomponent synthetic approach. The reaction involves a series of condensation and dehydrogenation phases that produce a particular C-C and C-N bond configuration. Deoxygenating of the alcohols is accomplished through condensation, whereas aromatization is achieved via dehydrogenations. This sustainable multicomponent synthesis is catalyzed by PN5P-Ir-pincer complexes with high efficiency (Deibl N, *et. al.*, 2015).




Fig. 10: Synthesis via Viable multicomponent synthesis

- **Synthesis of 2-substituted pyrimidine-5-carboxylic ester:** A simple and convenient method for the synthesis of several 2-substituted pyrimidine-5-carboxylic esters has been reported. In this approach, the sodium salt of 3,3-dimethoxy-2-methoxycarbonylpropen-1-ol is treated with various amidinium salts to produce the target compounds (Zhichkin P, *et. al.*, 2002).




Fig. 11: Synthesis of 2-substituted pyrimidine-5-carboxylic ester

- **Synthesis of densely substituted pyrimidines:** Synthesis of C4-heteroatom derivatized pyrimidines is accomplished by condensation of cyanic acid analogues with N-vinyl/aryl amides. In this reaction, the utilization of cyanic bromide and thiocyanatomethane offers flexible azaheterocycles poised for additional substitution (Ahmad OK, *et. al.*, 2009) and (Gayon E, *et. al.*, 2012).




Fig. 12: Synthesis of densely substituted pyrimidines

- **Synthesis via stereo selective access to beta-enaminones under NaOH catalysis:** This strategy consists of a rearrangement of propargylic hydroxylamines under NaOH catalysis, which permits efficient and stereo selective access to Cbz-protected beta-enaminones. Subsequent preparation of pyrimidines displays the synthetic application of these beta-enaminones (Gao F, *et. al.*, 2019).




Fig. 13: Synthesis via stereo selective access to beta-enaminones under NaOH catalysis

3. Structure Activity Relationship (SAR) of Pyrimidine:

Pyrimidine analogue SAR studies investigate the link between chemical structure and biological activity (Xiao J, *et. al.*, 2011). Pyrimidine ring modifications can also affect pharmacokinetic properties, such as solubility, permeability, and metabolic stability. Medicinal chemists use SAR to design and synthesize pyrimidine analogues with improved properties, including potency, selectivity, and pharmacokinetics, for use as pharmaceuticals (Zhang Y, *et. al.*, 2017).




Fig. 14: Structure Activity Relationship (SAR) analysis of Pyrimidines

Here are some possible substitutions which can be done to alter the activities of pyrimidine such as solubility, permeability, and metabolic stability:

- The introduction of electron-releasing moieties at position-2 of the core pyrimidine in tetrahydrobenzo [4,5] thieno [2,3-d] pyrimidine causes activity enhancement (Fatahala SS, *et. al.*, 2017).
- The incorporation of a 4-methoxyphenyl moiety at position-7 of pyrrolo[2,3-d]pyrimidine results in activity enhancement (Abdelgawad MA, *et. al.*, 2018).
- Pyridodipyrimidine derivatives with electron-releasing moieties show superior anti-inflammatory potential compared to derivatives of the same type with electron-withdrawing moieties (Raffa D, *et. al.*, 2009).
- Both monoaryl- and bisaryl-substituted 2-amino-pyrimidines exhibit improved activities regardless of the length of the C-5 substituents in polysubstituted pyrimidines (Chang C, *et. al.*, 2021).

- The introduction of two aryl moieties at neighbouring positions-2 and -3 in imidazo [1,2-a]pyrimidines leads to activity enhancement (ur Rashid H, *et. al.*, 2021).
- The presence of a five-membered thiophene moiety at positions-4,6 and an electron-releasing chlorine atom at position-2 of tricyclic pyrimidines improve the anti-inflammatory potential of the target derivatives (Bruni F, *et. al.*, 1993).
- Substitution of the phenyl ring with a thien-2-yl moiety at position-2 in pyrazolo [1,5-a]pyrimidin-7-ones results in substantial activity enhancement (Nofal ZM, *et. al.*, 2011).
- The introduction of 3-chlorophenyl and carboxamide groups at positions-1 and -4 of the pyrazolone ring, respectively, causes substantial enhancement in anti-inflammatory activities of pyrazolediazenyl pyrimidines (Bakr RB, *et. al.*, 2016).
- Pyrimidines bearing a pyrazolyl group in a hybrid configuration with the pyrazolo [3,4-d]pyrimidine moiety exhibit significant anti-inflammatory activities (Almansa C, *et. al.*, 2001).
- Pyrazolo [1, 5-a]pyrimidines with a 6,7-arrangement show the best anti-inflammatory activities (Amr AE, *et. al.*, 2017).
- Pyrimidines containing cycloheptenes fused to their rings display better activities than derivatives containing cyclohexenes fused to their rings (Zarghi A, *et. al.*, 2011).
- The introduction of p-sulfonamide or p-methylsulfone on one of the two aromatic rings in diarylheterocycle encompassing pyrimidines contributes to activity enhancement (Reddy DS, *et. al.*, 2021).

4. Pharmaceutical Applications:

Pyrimidine analogues have found extensive applications in the field of pharmaceuticals due to their broad range of biological activities, making them promising candidates for drug discovery and development (Andrei G, *et. al.*, 2012).

Some of the pharmaceutical applications of pyrimidine analogue is depicted in Fig. 5 and include:

- **Anticancer agents:**

Pyrimidine analogues such as 5-fluorouracil, cytarabine, and gemcitabine are commonly used as chemotherapy drugs due to their ability to inhibit DNA synthesis and cell proliferation, making them effective against a variety of cancers (Laev SS, *et. al.*, 2015).

- **Antiviral agents:**

Pyrimidine analogues like acyclovir and valacyclovir are used as antiviral agents due to their ability to inhibit DNA polymerase, making them effective against viral infections such as herpes and varicella-zoster (Singh K, *et. al.*, 2009).

- **Anti-inflammatory agents:**

Pyrimidine analogues such as leflunomide and teriflunomide have been developed as anti-inflammatory drugs, as they inhibit the production of pro-inflammatory cytokines and leukotrienes (Capasso C., *et. al.*, 2014).

- **Antihypertensive agents:**

Pyrimidine analogues like amlodipine and nifedipine are calcium channel blockers used as antihypertensive agents, as they relax blood vessels and lower blood pressure (Matos LH, *et. al.*, 2018).

- **Antimicrobial agents:**

Pyrimidine analogues such as trimethoprim and sulfamethoxazole are used as antimicrobial agents due to their ability to inhibit bacterial dihydrofolate reductase, an enzyme essential for the synthesis of folic acid (Hawser S, *et. al.*, 2006).

CONCLUSION:

In conclusion, pyrimidine and its analogous are important heterocyclic compounds with various applications in chemistry and pharmacy. Pyrimidine is a fundamental building block of DNA and RNA and plays a critical role in cellular metabolism. Pyrimidine derivatives and analogous have been found to exhibit diverse biological activities, including antiviral, antibacterial, anticancer, and anti-inflammatory properties. The synthesis of pyrimidine can be achieved using various classical and modern methods. Pyrimidine analogous can be classified into pyrimidine nucleosides and pyrimidine derivatives, with the latter having

different substituents attached to the ring, which can significantly alter their biological activity. The significance of substituted pyrimidines and purines in living organisms makes them extensively studied by chemists. The development of pyrimidine-based drugs has led to the discovery of many important pharmaceuticals, including anticancer agents and antiviral drugs. Overall, the importance of pyrimidine and its analogous in various fields underscores the need for continued research in this area.

REFERENCES:

1. Obaid RJ, Mughal EU, Naeem N, Al-Rooqi MM, Sadiq A, Jassas RS, Moussa Z, Ahmed SA. Pharmacological significance of nitrogen-containing five and six-membered heterocyclic scaffolds as potent cholinesterase inhibitors for drug discovery. *Process Biochemistry*. 2022 Jun 11.
2. Mukherjee AK, Basu S, Sarkar N, Ghosh AC. Advances in cancer therapy with plant based natural products. *Current medicinal chemistry*. 2001 Oct 1;8(12):1467-86.
3. Jain KS, Arya N, Inamdar NN, Auti PB, Unawane SA, Puranik HH, Sanap MS, Inamke AD, Mahale VJ, Prajapati CS, Shishoo CJ. The chemistry and bio-medicinal significance of pyrimidines & condensed pyrimidines. *Curr Top Med Chem*. 2016 Jan 1;16(28):3133-74.
4. Pels Rijcken WR, Hooghwinkel GJ, Ferwerda W. Pyrimidine metabolism and sugar nucleotide synthesis in rat liver. *Biochemical Journal*. 1990 Mar 15;266(3):777-83.
5. Choy N, Blanco B, Wen J, Krishan A, Russell KC. Photochemical and thermal Bergman cyclization of a pyrimidine enediynol and enediynone. *Organic Letters*. 2000 Nov 30;2(24):3761-4.
6. Stierne P, Debuigne A. Imine-based multicomponent polymerization: Concepts, structural diversity and applications. *Progress in Polymer Science*. 2022 Mar 2;101528.
7. Chiacchio MA, Iannazzo D, Romeo R, Giofrè SV, Legnani L. Pyridine and pyrimidine derivatives as privileged scaffolds in biologically active agents. *Current medicinal chemistry*. 2019 Dec 1;26(40):7166-95.
8. Lagoja IM. Pyrimidine as constituent of natural biologically active compounds. *Chemistry & Biodiversity*. 2005 Jan;2(1):1-50.
9. Keri RS, Hiremathad A, Budagumpi S, Nagaraja BM. Comprehensive review in current developments of benzimidazole-based medicinal chemistry. *Chemical biology & drug design*. 2015 Jul;86(1):19-65.
10. Selvam TP, James CR, Dniandev PV, Valzita SK. A mini review of pyrimidine and fused pyrimidine marketed drugs. *Research in Pharmacy*. 2015 Oct 29;2(4).
11. ur Rashid H, Martines MA, Duarte AP, Jorge J, Rasool S, Muhammad R, Ahmad N, Umar MN. Research developments in the syntheses, anti-inflammatory activities and structure–activity relationships of pyrimidines. *RSC advances*. 2021;11(11):6060-98.
12. Grazia Martina M, Giannessi L, Radi M. Multicomponent Synthesis of Purines and Pyrimidines: From the Origin of Life to New Sustainable Approaches for Drug-Discovery Applications. *European Journal of Organic Chemistry*. 2023 Jan 10;26(2):e202201288.
13. Gangjee A, Zeng Y, McGuire JJ, Mehraein F, Kisliuk RL. Synthesis of classical, three-carbon-bridged 5-substituted furo [2, 3-d] pyrimidine and 6-substituted pyrrolo [2, 3-d] pyrimidine analogues as antifolates. *Journal of medicinal chemistry*. 2004 Dec 30;47(27):6893-901.
14. Polshettiwar V, Varma RS. Microwave-assisted organic synthesis and transformations using benign reaction media. *Accounts of chemical research*. 2008 May 20;41(5):629-39.
15. Pałasz A, Cieź D. In search of uracil derivatives as bioactive agents. Uracils and fused uracils: Synthesis, biological activity and applications. *European Journal of Medicinal Chemistry*. 2015 Jun 5;97:582-611.
16. Sasada T, Kobayashi F, Sakai N, Konakahara T. An unprecedented approach to 4, 5-disubstituted pyrimidine derivatives by a ZnCl₂-catalyzed three-component coupling reaction. *Organic letters*. 2009 May 21;11(10):2161-4.
17. Foley DJ, Waldmann H. Ketones as strategic building blocks for the synthesis of natural product-inspired compounds. *Chemical Society Reviews*. 2022.

18. Jadhav SD, Singh A. Oxidative annulations involving DMSO and formamide: K₂S₂O₈ mediated syntheses of quinolines and pyrimidines. *Organic letters*. 2017 Oct 20;19(20):5673-6.
19. Su L, Sun K, Pan N, Liu L, Sun M, Dong J, Zhou Y, Yin SF. Cyclization of ketones with nitriles under base: A general and economical synthesis of pyrimidines. *Organic letters*. 2018 May 23;20(11):3399-402.
20. Chu XQ, Cao WB, Xu XP, Ji SJ. Iron catalysis for modular pyrimidine synthesis through β -ammoniation/cyclization of saturated carbonyl compounds with amidines. *The Journal of Organic Chemistry*. 2017 Jan 20;82(2):1145-54.
21. Guo W, Li C, Liao J, Ji F, Liu D, Wu W, Jiang H. Transition metal free intermolecular direct oxidative C–N bond formation to polysubstituted pyrimidines using molecular oxygen as the sole oxidant. *The Journal of Organic Chemistry*. 2016 Jul 1;81(13):5538-46.
22. Zhan JL, Wu MW, Chen F, Han B. Cu-catalyzed [3+ 3] annulation for the synthesis of pyrimidines via β -C (sp³)–H functionalization of saturated ketones. *The Journal of Organic Chemistry*. 2016 Dec 2;81(23):11994-2000.
23. Deibl N, Ament K, Kempe R. A sustainable multicomponent pyrimidine synthesis. *Journal of the American Chemical Society*. 2015 Oct 14;137(40):12804-7.
24. Zhichkin P, Fairfax DJ, Eisenbeis SA. A general procedure for the synthesis of 2-substituted pyrimidine-5-carboxylic esters. *Synthesis*. 2002;2002(06):0720-2.
25. Hill MD, Movassaghi M. Direct synthesis of substituted pyrimidines and quinazolines. *Synthesis*. 2008 Mar;2008(05):823-7.
26. Ahmad OK, Hill MD, Movassaghi M. Synthesis of densely substituted pyrimidine derivatives. *The Journal of Organic Chemistry*. 2009 Nov 6;74(21):8460-3.
27. Gayon E, Szymczyk M, Gérard H, Vrancken E, Campagne JM. Stereoselective and catalytic access to β -enaminones: An entry to pyrimidines. *The Journal of Organic Chemistry*. 2012 Oct 19;77(20):9205-20.
28. Gao F, Wang T, Xiao J, Huang G. Antibacterial activity study of 1, 2, 4-triazole derivatives. *European journal of medicinal chemistry*. 2019 Jul 1;173:274-81.
29. Xiao J, Marugan JJ, Zheng W, Titus S, Southall N, Cherry JJ, Evans M, Androphy EJ, Austin CP. Discovery, synthesis, and biological evaluation of novel SMN protein modulators. *Journal of medicinal chemistry*. 2011 Sep 22;54(18):6215-33.
30. Zhang Y, Luo L, Han C, Lv H, Chen D, Shen G, Wu K, Pan S, Ye F. Design, synthesis, and biological activity of tetrahydrobenzo [4, 5] thieno [2, 3-d] pyrimidine derivatives as anti-inflammatory agents. *Molecules*. 2017 Nov 13;22(11):1960.
31. Fatahala SS, Mohamed MS, Youns M, Abd-El Hameed RH. Synthesis and evaluation of cytotoxic activity of some pyrroles and fused pyrroles. *Anti-Cancer Agents in Medicinal Chemistry (Formerly Current Medicinal Chemistry-Anti-Cancer Agents)*. 2017 Jun 1;17(7):1014-25.
32. Abdelgawad MA, Bakr RB, Azouz AA. Novel pyrimidine-pyridine hybrids: synthesis, cyclooxygenase inhibition, anti-inflammatory activity and ulcerogenic liability. *Bioorganic Chemistry*. 2018 Apr 1;77:339-48.
33. Raffa D, Maggio B, Plescia F, Cascioferro S, Raimondi MV, Plescia S, Cusimano MG. Pyrazolo [3, 4-d] pyrimidine Derivatives as COX-2 Selective Inhibitors: Synthesis and Molecular Modelling Studies. *Archiv der Pharmazie: An International Journal Pharmaceutical and Medicinal Chemistry*. 2009 Jun;342(6):321-6.
34. Chang C, Chen W, Chen Y, Chen Y, Chen Y, Ding F, Fan C, Fan HJ, Fan Z, Gong C, Gong Y. Recent progress on two-dimensional materials. *Acta Phys.-Chim. Sin.* 2021 Nov 15;37(12):2108017.
35. ur Rashid H, Martines MA, Duarte AP, Jorge J, Rasool S, Muhammad R, Ahmad N, Umar MN. Research developments in the syntheses, anti-inflammatory activities and structure–activity relationships of pyrimidines. *RSC advances*. 2021;11(11):6060-98.
36. Bruni F, Costanzo A, Selleri S, Guerrini G, Fantozzi R, Pirisino R, Brunelleschi S. Synthesis and study of the anti-inflammatory properties of some pyrazolo [1, 5-a] pyrimidine derivatives. *Journal of pharmaceutical sciences*. 1993 May 1;82(5):480-6.
37. Nofal ZM, Fahmy HH, Zarea ES, El-Eraky W. Synthesis of new pyrimidine derivatives with evaluation of their anti-inflammatory and analgesic activities. *Acta Pol Pharm*. 2011 Jul 1;68(4):507-17.
38. Bakr RB, Azouz AA, Abdellatif KR. Synthesis, cyclooxygenase inhibition, anti-inflammatory evaluation and ulcerogenic liability of new 1-phenylpyrazolo [3, 4-d] pyrimidine derivatives. *Journal of enzyme inhibition and medicinal chemistry*. 2016 Nov 2;31(sup2):6-12.

39. Almansa C, de Arriba AF, Cavalcanti FL, Gómez LA, Miralles A, Merlos M, García-Rafanell J, Forn J. Synthesis and SAR of a new series of COX-2-selective inhibitors: pyrazolo [1, 5-a] pyrimidines. *Journal of medicinal chemistry*. 2001 Feb 1;44(3):350-61.
40. Amr AE, Abdalla MM, Ouf NH. Anti-Angiogenic Effects of Some (5-hydroxy-4-methyl-2-((1-methyl-1H-indol-3-yl) methyl)-Thieno [2, 3-d] Pyrimidine Derivatives. *Journal of Computational and Theoretical Nanoscience*. 2017 Jan 1;14(1):448-53.
41. Zarghi A, Arfaei S. Selective COX-2 inhibitors: a review of their structure-activity relationships. *Iranian journal of pharmaceutical research: IJPR*. 2011;10(4):655.
42. Reddy DS, Kongot M, Kumar A. Coumarin hybrid derivatives as promising leads to treat tuberculosis: Recent developments and critical aspects of structural design to exhibit anti-tubercular activity. *Tuberculosis*. 2021 Mar 1;127:102050.
43. Galmarini CM, Mackey JR, Dumontet C. Nucleoside analogues and nucleobases in cancer treatment. *The lancet oncology*. 2002 Jul 1;3(7):415-24.
44. Andrei G, Topalis D, Fiten P, McGuigan C, Balzarini J, Opdenakker G, Snoeck R. In vitro-selected drug-resistant varicella-zoster virus mutants in the thymidine kinase and DNA polymerase genes yield novel phenotype-genotype associations and highlight differences between antiherpesvirus drugs. *Journal of virology*. 2012 Mar 1;86(5):2641-52.
45. Laev SS, Salakhutdinov NF. Anti-arthritis agents: progress and potential. *Bioorganic & medicinal chemistry*. 2015 Jul 1;23(13):3059-80.
46. Singh K, Arora D, Singh K, Singh S. Genesis of dihydropyrimidinone calcium channel blockers: recent progress in structure-activity relationships and other effects. *Mini reviews in medicinal chemistry*. 2009 Jan 1;9(1):95.
47. Capasso C, Supuran CT. Sulfa and trimethoprim-like drugs—antimetabolites acting as carbonic anhydrase, dihydropteroate synthase and dihydrofolate reductase inhibitors. *Journal of enzyme inhibition and medicinal chemistry*. 2014 Jun 1;29(3):379-87.

