



## A Critical Review of Derivative Ultraviolet Spectrophotometry in Pharmaceutical Analysis

Pratiksha S. Karwande<sup>1\*</sup>, Bhakti P. Gunjal<sup>1</sup>, Shashank S. Morkile<sup>1</sup>, Vijaya U. Barge<sup>2</sup>

<sup>1</sup>Department of Pharmaceutical Quality Assurance, Shankarrao Ursal College of Pharmaceutical Sciences & Research Centre, Kharadi, Pune – 411014. India.

<sup>2</sup>Vice Principal & Assistant Professor, Department of Pharmaceutical Quality Assurance, Shankarrao Ursal College of Pharmaceutical Sciences & Research Centre, Kharadi, Pune – 411014. India.

Received: 25 January 2026

Revised: 10 February 2026

Accepted: 26 February 2026

### ABSTRACT

Derivative UV spectrophotometry is an advanced analytical technique widely used in pharmaceutical analysis for qualitative and quantitative determination of drugs. The method enhances spectral resolution and improves analytical sensitivity compared to conventional UV spectroscopy. By transforming the normal absorbance spectrum into its derivative form, the method allows the precise estimation of analytes in complex mixtures without prior separation. Derivative spectrophotometry has gained significant importance in pharmaceutical quality control due to its simplicity, cost-effectiveness, and rapid analysis capability. This review focuses on the theoretical principles, instrumentation, mathematical background, and pharmaceutical applications of derivative UV spectroscopy. Additionally, advantages, limitations, and recent developments in derivative spectroscopic techniques are discussed. The technique is particularly useful for the simultaneous estimation of drugs in multicomponent formulations, improving accuracy and selectivity. Thus, derivative UV spectrophotometry serves as an effective analytical tool in pharmaceutical research, method development, and quality control.

**Keywords:** Derivative spectroscopy, Ultraviolet spectroscopy, Pharmaceutical analysis, Spectral resolution, Analytical method development, Quality control, Drug estimation.

### 1. INTRODUCTION

Ultraviolet-visible (UV-Vis) spectroscopy is one of the most commonly used analytical techniques in pharmaceutical analysis<sup>[16,17]</sup>. It is widely applied for drug identification, purity testing, and quantitative estimation in bulk drugs and pharmaceutical formulations. The technique is based on the absorption of ultraviolet or visible radiation by molecules, resulting in electronic transitions between molecular orbitals<sup>[16]</sup>.

Despite its widespread use, conventional UV spectrophotometry faces certain limitations during the analysis of multicomponent pharmaceutical formulations. Many drugs exhibit overlapping absorption spectra, which makes their simultaneous determination difficult using normal spectrophotometric methods<sup>(1,3)</sup>. Spectral interference between analytes often reduces the accuracy and selectivity of the analytical method. To overcome these challenges, several advanced spectroscopic approaches have been developed, among which derivative ultraviolet spectrophotometry has gained significant attention<sup>[1]</sup>.

Derivative spectrophotometry is a mathematical transformation technique in which the normal absorption spectrum is converted into its derivative forms. The derivative spectra represent the rate of change of absorbance with respect to wavelength<sup>[3]</sup>. This transformation enhances spectral resolution, enabling the detection of small spectral features that may not be visible in the conventional zero-order spectrum<sup>[3,4]</sup>. As a result, derivative spectrophotometry allows accurate determination of analytes even in the presence of overlapping spectral bands.

The development of modern computerised UV spectrophotometers has further increased the applicability of derivative spectrophotometric methods in pharmaceutical analysis<sup>[7]</sup>. Advanced software tools enable rapid calculation of first-order, second-order, and higher-order derivative spectra with improved precision and reduced noise interference<sup>[3,7]</sup>. These developments have made derivative spectrophotometry a valuable analytical tool for pharmaceutical scientists.

In pharmaceutical research and quality control laboratories, derivative UV spectrophotometry has been successfully applied for simultaneous estimation of drugs in combined dosage forms, impurity detection, stability studies, and dissolution testing<sup>[7,9]</sup>. The method offers several advantages, including improved selectivity, enhanced sensitivity, minimal sample preparation, and cost-effective analysis compared with other advanced analytical techniques such as high-performance liquid chromatography (HPLC)<sup>[16]</sup>.

Because of these advantages, derivative spectrophotometry has become an important analytical technique in pharmaceutical method development and routine quality control analysis. The present review aims to provide a comprehensive overview of the principles, instrumentation, and applications of derivative ultraviolet spectrophotometry in pharmaceutical analysis.

This review aims to provide a comprehensive overview of the principles, instrumentation, types, and pharmaceutical applications of derivative ultraviolet spectrophotometry, along with its advantages, limitations, and recent advancements in analytical techniques.

## 2. INSTRUMENTATION OF UV-VISIBLE SPECTROPHOTOMETER

A UV-visible spectrophotometer is an analytical instrument used to measure the absorption of ultraviolet and visible radiation by a sample<sup>[16]</sup>. The instrument consists of several important components that work together to produce an absorption spectrum and determine the concentration of pharmaceutical compounds.


### Main components :

- **Light source :**

Provides ultraviolet and visible radiation. A deuterium lamp is commonly used for the UV region ( 190-380 nm), and a tungsten lamp is used for the visible region ( 380-800 nm)<sup>[16]</sup>.

- **Monochromator :**

It selects a specific wavelength of light from the polychromatic radiation produced by the light source. It usually consists of a prism or diffraction grating<sup>[21]</sup>.



*Figure 1. Instrumentation of UV Spectrophotometry*

- **Sample holder ( cuvette) :**

The sample solution is placed in a quartz cuvette for UV analysis because quartz allows UV radiation to pass through.

- **Detector :**

The detector measures the intensity of transmitted light and converts it into an electrical signal. Common detectors include photodiodes and photomultiplier tubes<sup>[16]</sup>.



- **Amplifier and readout system :**

The electrical signal produced by the detector is amplified and displayed as absorbance or transmittance on a digital screen or computer.

### 3. PRINCIPLE OF DERIVATIVE SPECTROPHOTOMETRY

Derivative spectrophotometry is a mathematical transformation of the normal absorption spectrum ( zero-order spectrum) into its derivative forms with respect to wavelength<sup>[2]</sup>. In this technique, the derivative of absorbance is measured instead of the direct absorbance value. The derivative spectra may be expressed as first-order, second-order, or higher-order derivatives of the original spectrum.

The Mathematical expressions are:

$$\text{First derivative} : D^1 = dA/d\lambda$$

$$\text{Second derivative} : D^2 = d^2A/d\lambda^2$$

$$\text{Third derivative} : D^3 = d^3A/d\lambda^3$$

Where A is absorbance, and  $\lambda$  is wavelength

The derivative spectrum represents the rate of change of absorbance with wavelength, which improves spectral resolution and allows better separation of overlapping absorption bands<sup>[3]</sup>. Therefore, derivative spectrophotometry is widely used in pharmaceutical analysis for the simultaneous estimation of drugs in multicomponent formulations and for reducing interference from excipients and baseline shifts<sup>[1]</sup>.

### 4. TYPES OF DERIVATIVE SPECTROPHOTOMETRY

Derivative spectrophotometry is classified according to the order of the derivative of the absorption spectrum. The commonly used types include<sup>[2]</sup>:


**Table 1. Types of Derivative Spectrophotometry**

SR.NO	DERIVATIVE ORDER	SYMBOL	DESCRIPTION	ANALYTICAL SIGNIFICANCE
1.	Zero-order spectrum	D <sup>0</sup>	Represents the normal absorption spectrum obtained by plotting absorbance against wavelength in conventional UV-visible spectrophotometry.	Commonly used for routine qualitative and quantitative determination of pharmaceutical compounds.
2.	First-order derivative	D <sup>1</sup>	Corresponds to the first derivative of absorbance with respect to wavelength(dA/dλ), representing the rate of change of absorbance with wavelength.	Enhances spectral resolution and is widely used for the analysis of drugs with overlapping absorption spectra.
3.	Second-order derivative	D <sup>2</sup>	Represents the second derivative of absorbance with respect to wavelength (d <sup>2</sup> A/dλ <sup>2</sup> ), providing sharper spectral features and improved discrimination of closely spaced peaks.	Applied for improved selectivity in multicomponent pharmaceutical formulations.
4.	Third/Higher-order derivatives	D <sup>3</sup> , D <sup>4</sup>	Third-order and higher derivatives were obtained through further mathematical differentiation of the absorption spectrum.	Provide additional spectral information, but they are less frequently used due to amplification of spectral noise.

### 5. DERIVATIVE SPECTRA IN UV SPECTROPHOTOMETRY

The derivative spectra obtained in UV spectrophotometry are shown below in order of differentiation.

## 5.1 ZERO-ORDER SPECTRUM



**Figure 2 Zero-Order Spectrum**

The zero-order spectrum represents the normal UV absorption spectrum obtained in UV-visible spectrophotometry. In this spectrum, absorbance (A) is plotted against wavelength.

X-axis: Wavelength (nm)

Y-axis: Absorbance


The highest point of the curve is called  $\lambda_{\max}$  (maximum absorbance). Quantitative analysis of drugs in conventional UV spectroscopy is generally performed at  $\lambda_{\max}$  because it provides maximum sensitivity and accuracy<sup>[16]</sup>.

### Importance :

- 1) Determination of the wavelength of maximum absorbance ( $\lambda_{\max}$ ).
- 2) Provide maximum sensitivity for quantitative analysis.
- 3) Serve as the fundamental spectrum for deriving derivative spectra.
- 4) Assist in UV spectrophotometric method development.

## 5.2 FIRST-ORDER DERIVATIVE SPECTRUM :

The first-order derivative spectrum is obtained by mathematically differentiating the normal UV absorption spectrum. In this spectrum, the rate of change of absorbance with respect to wavelength is plotted.



*Figure 2. First-Order Derivative Spectrum*

Mathematically :

$$dA/d\lambda$$

where

A = absorbance

$\lambda$  = wavelength

The spectrum shows positive and negative peaks.

The point where the curve crosses the baseline is called the zero-crossing point.


The original  $\lambda_{\max}$  of the zero spectrum corresponds to the zero-crossing point in the first-order spectrum<sup>[3]</sup>.

#### **Importance :**

First-order derivative spectroscopy helps to :

- 1) Improve spectral resolution.
- 2) Reduce baseline interference.
- 3) Allows simultaneous estimation of drugs in mixtures.

### 5.3 SECOND-ORDER DERIVATIVE SPECTRUM :



*Figure 3. Second-Order Derivative Spectrum*

The second-order derivative spectrum represents the second derivative of absorbance with respect to wavelength.

Which is mathematically expressed as :

$$d^2A/d\lambda^2$$

where,

A= Absorbance

λ= wavelength


The second-order derivative spectrum produces sharper peaks and better resolution, allowing separation of closely overlapping spectral bands<sup>[2]</sup>.

#### **Importance :**

- 1) Improve spectral resolution between overlapping peaks.
- 2) Enhance peak discrimination in complex mixtures.
- 3) Reduce interference from neighboring absorption bands.
- 4) Provides better selectivity in pharmaceutical analysis.

#### 5.4 THIRD-ORDER DERIVATIVE SPECTRUM

The Third-order derivative spectrum represents the third derivative of absorbance with respect to wavelength.



**Figure 4. Third-Order Derivative Spectrum**

Which is mathematically expressed as:

$$d^3A / d\lambda^3$$

where,

A= absorbance

$\lambda$  = wavelength

The third-order derivative spectrum provides very high spectral resolution, which helps in distinguishing closely overlapping absorption bands. However, higher-order derivatives are less commonly used because they may increase spectral noise<sup>[35]</sup>.

#### **Importance :**

- 1) Provides higher spectral resolution.
- 2) Distinguishing overlapping peaks very closely.
- 3) Improve selectivity in complex mixtures.
- 4) Enhance analytical accuracy in multicomponent analysis.

Spectrophotometry remains an efficient and reliable analytical tool when appropriate instrumental parameters are carefully optimised. With ongoing advancements in spectroscopic instrumentation and data processing techniques, the application of derivative spectrophotometry is expected to expand further in pharmaceutical research, quality control, and analytical method development.

#### **6. GENERAL ANALYTICAL PROCEDURE OF DERIVATIVE UV SPECTROPHOTOMETRY**

The general procedure for derivative UV spectrophotometric analysis involves several steps for the accurate determination of pharmaceutical compounds<sup>[1,3]</sup>.



### **Step 1: Preparation of standard solution**

An accurately weighed quantity of the drug is dissolved in a suitable solvent such as methanol, ethanol, or distilled water to prepare a stock solution of known concentration.

### **Step 2: Preparation of working standards**

The stock solution is further diluted with the selected solvent to obtain working solutions within the required concentration range for analysis.

### **Step 3: Recording the zero-order spectrum**

The prepared solution is scanned using a UV-visible spectrophotometer in the wavelength range of 200-400 nm to obtain the normal absorption spectrum (zero-order spectrum)<sup>[2,3]</sup>.

### **Step 4: Conversion to derivative spectrum**

Using the spectrophotometer software, the recorded spectrum is mathematically converted into first-order, second-order, and higher-order derivative spectra.

### **Step 5: Selection of analytical wavelength**

A suitable peak amplitude or zero-crossing wavelength is selected from the derivative spectrum for quantitative analysis.

### **Step 6: Calibration curve preparation**

Derivative absorbance values are plotted against concentration to construct a calibration curve.

### **Step 7: Sample analysis**

The derivative absorbance of the sample solution is measured, and the concentration of the drug is calculated using the calibration equation.

This procedure allows accurate determination of drugs even in the presence of overlapping spectra, making derivative spectrophotometry particularly useful for multicomponent formulations<sup>[3]</sup>.

## **7. MATERIALS AND METHOD**

The present review is based on the systematic collection and analysis of previously published literature related to derivative ultraviolet spectrophotometry. Relevant research articles, review papers, and standard textbooks were referred from scientific databases such as Google Scholar, PubMed, and ScienceDirect.

A UV-visible spectrophotometer equipped with appropriate software for derivative spectral analysis was considered as the primary analytical instrument. Common solvents such as methanol, ethanol, and distilled water were used as reference media in reported studies.

The collected data were critically analysed to understand the principles, instrumentation, derivative orders, and pharmaceutical applications of derivative UV spectrophotometry.

## **8. RESULTS AND DISCUSSION**

Derivative ultraviolet spectrophotometry is an effective analytical technique that significantly improves spectral resolution and enables accurate analysis of drugs in complex mixtures. The transformation of zero-order spectra into higher-order derivatives allows better separation of overlapping peaks, which is a major limitation in conventional UV spectrophotometry.

The technique is widely applied in pharmaceutical analysis for the simultaneous estimation of multicomponent formulations, impurity detection, and stability studies. Compared to chromatographic techniques such as HPLC, derivative UV spectrophotometry



is simpler, more economical, and requires less time for analysis. However, it may show lower sensitivity compared to advanced chromatographic methods.

Overall, derivative spectrophotometry provides a reliable and cost-effective alternative for routine quality control analysis in the pharmaceutical industry.

## **9. APPLICATIONS IN PHARMACEUTICAL ANALYSIS.**

Derivative spectrophotometry has numerous applications in pharmaceutical analysis due to its ability to improve spectral resolution and analytical selectivity<sup>[1,3]</sup>.

### **1) Quantitative determination of drugs :**

Derivative spectrophotometry is widely used for the quantitative analysis of drugs in bulk materials and pharmaceutical dosage forms.

### **2) Simultaneous estimation of multicomponent formulations:**

It helps in the simultaneous determination of two or more drugs in combination formulations without prior separation.

### **3) Resolution of overlapping spectra :**

The technique improves spectral resolution and allows accurate determination of drugs with overlapping absorption spectra.

### **4) Quality control analysis:**

It is commonly used in pharmaceutical industries for routine quality control of drug products.

### **5) Stability studies :**

Derivative spectroscopy can detect small spectral changes and is useful for studying drug degradation and stability.

### **6) Detection of impurities and trace analysis :**

It can be used to identify and quantify impurities present in pharmaceutical formulations.

### **7) Analysis in biological samples :**

The method is also applied for the determination of drugs in biological fluids such as plasma and urine.

### **8) Determination of drugs in combination dosage form :**

Derivative spectrophotometry is useful for determining the concentration of drugs present in combination pharmaceutical formulations.

### **9) Method development and validation :**

Derivative spectrophotometry is applied in the development and validation of analytical methods for accurate drug estimation.

### **10) Dissolution testing analysis :**

The technique can be used to analyse drug release during dissolution studies of pharmaceutical dosage forms.

### **11) Analysis of herbal and natural products :**

The technique is applied for the determination of active pharmaceutical ingredients present in herbal formulations and plant extracts.



## **12) Environmental analysis of pharmaceutical residues :**

Derivative spectrophotometry is used to detect pharmaceutical compounds present in environmental samples such as wastewater and surface water.

## **13) Detection of drug degradation products :**

Derivative spectroscopy helps to identify small spectral changes caused by degradation products during forced degradation studies.

## **14) Chemometric-assisted derivative spectrophotometry :**

Derivative spectroscopy combined with chemometric techniques improves the analysis of complex pharmaceutical mixtures.

## **10. ADVANTAGES AND LIMITATIONS OF DERIVATIVE SPECTROPHOTOMETRY**

### **ADVANTAGES :**

- Derivative spectrophotometry offers several advantages over conventional UV spectrophotometric methods<sup>[2,3]</sup>.
- Improves spectral resolution and helps separate overlapping absorption bands.
- Reduces baseline drift and background interference.
- Enables simultaneous estimation of multiple drugs in mixture formulations.
- Requires simple instrumentation compared to chromatographic methods.
- Provides a rapid and cost-effective analytical technique for routine pharmaceutical analysis.

### **LIMITATIONS**

- Despite its advantages, derivative spectrophotometry also has certain limitations <sup>[1,2]</sup>.
- Higher-order derivatives may increase spectral noise.
- Requires careful selection of derivative order and wavelength.
- Sensitivity may be lower compared to chromatographic methods such as HPLC.
- Instrumental parameters must be properly optimised for accurate results.

## **11. FUTURE PERSPECTIVE OF DERIVATIVE SPECTROPHOTOMETRY**

- Development of hyphenated analytical techniques combining derivative spectrophotometry with advanced spectroscopic tools<sup>[1,3]</sup>.
- Application of artificial intelligence and machine learning for improved spectral data interpretation.
- Use of derivative spectrophotometry in real-time pharmaceutical process monitoring during drug manufacturing.
- Expansion of the technique for the analysis of complex herbal and natural product formulation.
- Increased use in environmental monitoring of pharmaceutical contaminants in water and soil samples.
- Integration with chemometric modelling to improve multicomponent drug analysis.



## 12. CONCLUSION

Derivative ultraviolet spectrophotometry has emerged as a powerful and versatile analytical technique in pharmaceutical analysis. By mathematically transforming the normal absorption spectrum into its derivative forms, the method significantly improves spectral resolution and enables the accurate determination of drugs even in the presence of overlapping spectra. These characteristics make derivative spectrophotometry particularly useful for the simultaneous estimation of multicomponent pharmaceutical formulations.

The technique has been widely applied in various areas of pharmaceutical analysis, including quantitative drug determination in bulk and dosage form, quality control analysis, stability studies, impurity detection, and analysis of drugs in biological samples. Compared with more complex analytical techniques such as chromatographic methods, derivative spectrophotometry offers advantages such as simplicity, rapid analysis, minimal sample preparation, and lower operational cost<sup>[1,3]</sup>.

Despite certain limitations, such as increased spectral noise at higher derivative orders, derivative spectrophotometry remains an efficient and reliable analytical tool when appropriate instrumental parameters are carefully optimised. With ongoing advancements in spectroscopic instrumentation and data processing techniques, the application of derivative spectrophotometry is expected to expand further in pharmaceutical research, quality control, and analytical method development.

## ACKNOWLEDGEMENT

The authors are grateful to the management of PDEA's Shankarrao Ursal College of Pharmaceutical Sciences and Research Centre, Kharadi, Pune, for providing the necessary facilities and support to carry out this review study. The authors express their sincere gratitude to Dr. Vijaya Barge for valuable guidance, continuous support, and encouragement throughout the preparation of this manuscript. The authors also acknowledge all the researchers and authors whose published work has been consulted and cited in this review article.

## REFERENCES

1. Redasani VK, Patel PR, Marathe DY, Chaudhari SR, Shirkhedkar AA, Surana SJ. A review on derivative UV spectrophotometry analysis of drugs in pharmaceutical formulations and biological samples. *J Chil Chem Soc.* 2018;63(3):4126-4134.
2. Karpinska J. Basic principles and analytical applications of derivative spectrophotometry. *Talanta.* 2001;64:801-822.
3. Rojas F, Ojeda C, Pavon J. Derivative spectrophotometry in analytical chemistry. *Talanta.* 1988;35:753-761.
4. Kus S, Marzenko Z, Obarski N. Derivative spectrophotometry in analytical chemistry. *Chem Anal.* 1996;41:899-927.
5. Salem H, Abdallah O. Application of derivative spectroscopy in pharmaceutical analysis. *AM J Appl Sci.* 2007;4:709-717.
6. Cameron D, Moffatt D. Application of derivative spectroscopy in analytical chemistry. *Appl Spectrosc.* 1987;41:539-544.
7. Koba M, Koba K, Przyborowski L. Application of derivative spectrophotometry in pharmaceutical analysis. *Acta Pol Pharm.* 2008;65:289-294.
8. Shah D, Bhatt K, Mehta R, Baldania S. Derivative spectrophotometric determination of pharmaceutical compounds in dosage forms. *J AOAC Int.* 2008;91:1075-1082.
9. Murillo J, Lernus J, Garcia L. Derivative spectrophotometric determination of drugs in pharmaceutical formulations. *J Pharma Biomed Anal.* 1996;14:257-266.
10. Singh DK, Verma R. Comparison of second derivative spectrophotometric and RP-HPLC methods for determination of prednisolone in pharmaceutical formulations. *Anal Sci.* 2007;23:1241-1243.
11. Lavanya G, Sunil M, Eswarudu M, Naga Raji P. UV spectrophotometric methods for pharmaceutical analysis. *Int J Pharm Sci Rev Res.* 2013.
12. Upstone S. Ultraviolet- visible absorption spectrophotometry in analytical chemistry. *Encyclopedia of Analytical Chemistry.*
13. Marzenko Z. Spectrophotometric determination of elements. Wiley.
14. Snyder LR, Kirkland JJ. Introduction to modern liquid chromatography. Wiley.
15. Harris DC. Quantitative chemical analysis. Freeman Publishers.
16. Skoog DA, Holler FJ, Crouch SR. Principles of Instrumental Analysis. 6<sup>th</sup> ed.
17. Skoog DA, West DM, Holler FJ. Fundamentals of Analytical Chemistry.
18. Christian GD. Analytical Chemistry. 6<sup>th</sup> ed. Wiley.
19. Harvey D. Modern Analytical Chemistry. McGraw-hill.
20. Miller JN, Miller JC. Statistics and Chemometrics for Analytical Chemistry. Pearson.
21. Willard HH, Merritt LL, Dean JA, Settle FA. Instrumental Methods of Analysis. CBS Publishers.
22. Rouessac F, Rouessac A. Chemical Analysis: Modern Instrumentation Methods and Techniques.
23. Beckett AH, Stenlake JB. Practical Pharmaceutical Chemistry. CBS Publishers.
24. Chatwal GR, Anand SK. Instrumental Methods of Chemical Analysis. Himalaya Publishing House.
25. Sharma BK. Instrumental Methods of Chemical Analysis. Goel Publishing.



26. Sethi PD. Quantitative Analysis of Drugs in Pharmaceutical Formulations. CBS Publishers.
27. Jeffery GH, Bassett J, Mendham J, Denney RC. Vogel's Textbook of Quantitative Chemical Analysis.
28. ICH. Q2(f Analytical ProceduresR1): Validation of Analytical Procedures: Text and Methodology. 2005.
29. United States Pharmacopoeia (USP). Spectrophotometric Methods.
30. British Pharmacopoeia. Spectrophotometric Methods.
31. Kellner R, Mermet JM, Otto M. Analytical Chemistry: A Modern Approach to Analytical Science.
32. Harris DC. Exploring Chemical Analysis.
33. Rouessac F. Chemical Analysis: Modern Instrumentation Methods.
34. Douglas A Skoog. Application of instrumental analysis in pharmaceutical chemistry.
35. Talsky G. Derivative Spectrophotometry in Analytical Chemistry.

How to cite this article:

Pratiksha S. Karwande et al. *Ijppr.Human*, 2026; Vol. 32 (3): 429-440.

Conflict of Interest Statement: All authors have nothing else to disclose.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.