International Journal of Pharmacy and Pharmaceutical Research (IJPPR)
Volume 32, Issue 7, July 2026 ijppr.humanjournals.com ISSN: 2349-7203

Inflammation: A Comprehensive Overview of Pathogenesis, Diagnosis, and Treatment

Mr. Shantakumar, Harish S, Pravin P, Preetha B, Sindhu E, Teena Elisa Thomas

Department of Pharmacology, United College of Pharmacy, Periyanaickenpalayam, Coimbatore. India.

Received: 27 May 2026 Revised: 15 June 2026 Accepted: 28 June 2026

ABSTRACT

Inflammation is an immunological reaction that protects the body from toxic stimuli, infections, and tissue damage. Immune cells, blood vessels, and inflammatory mediators all work together in concert. Acute inflammation helps heal and restore homeostasis, but chronic inflammation contributes to the development of several illnesses, including cancer, diabetes, rheumatoid arthritis, inflammatory bowel disease, cardiovascular disorders, and neurological diseases. Important signalling pathways like NF-κB, MAPK, JAK/STAT, and COX control the inflammatory response. The mechanics of inflammation types, mediators, pathogenesis, diagnostic biomarkers, and treatment modalities are compiled in this chapter. It emphasizes recent developments in targeted therapies, natural bioactive compounds, experimental models for assessing anti-inflammatory activity, and traditional anti-inflammatory medications. To effectively prevent and treat inflammatory illnesses, it is imperative to comprehend these pathways. Objectives**:** To give a summary of the physiological relevance of the inflammatory process. To explain the many forms of inflammation, such as acute and chronic inflammation. To describe the molecular and cellular processes that underlie inflammation. To describe inflammation's pathophysiology, clinical signs, and diagnostic biomarkers. To examine how both natural and conventional anti-inflammatory drugs work.

Keywords: Inflammation, acute inflammation, chronic inflammation, anti-inflammatory agents, inflammatory mediators, natural products, and therapeutic strategies.

INTRODUCTION

Inflammation is the body's quick and general reaction to harmful things like germs, foreign objects, dead cells, irritants, or cancer cells. This natural immune response includes a teamwork of various immune cells, the walls of blood vessels, and important chemical signals that help protect the body. When you get hurt or sick, these components work together to fight off threats and begin the healing process [1]. Inflammation is a process in the body that can become more intense as people get older. When there are high levels of inflammatory markers, it can raise the chances of developing cardiovascular diseases (CVDs). These include conditions like high blood pressure (hypertension), the hardening of arteries (atherosclerosis), and even a fast decline in heart health leading to heart failure. This ongoing inflammation related to aging is often called "inflammaging [2].

Inflammation is the body's natural way of responding to harmful things like germs, injured cells, or toxic substances. This process involves various immune cells, blood vessels, and chemical signals that work together to fight off these dangers. When inflammation doesn't go away and becomes chronic, it can lead to the release of substances called cytokines that keep the inflammation going and create stress in the cells. An imbalance in the cell's energy factories, known as mitochondria, or in the cell's nucleus, can cause damage to DNA. When DNA gets damaged, the cell activates a special response system called the DNA Damage Response (DDR). This system is mainly controlled by proteins known as ATM and ATR kinases. These proteins change how genes are expressed and how the cell uses energy, which can ultimately lead to a state called senescence, where cells stop dividing and function less effectively 3.

Inflammation is a natural reaction that happens in our bodies when we encounter harmful things or situations. This includes when we get sick due to an infection, like a cold or the flu, or when we hurt ourselves, such as getting a cut or a bruise. Our body uses inflammation to help protect and heal itself 4.

TYPES OF INFLAMMATION

- ***Chronic inflammation*** is a long-lasting health issue that occurs when certain immune cells, like monocytes, macrophages, lymphocytes, and plasma cells, invade tissues. This condition can lead to damage in the affected areas and the formation of scar

tissue, known as fibrosis. One of the main problems with chronic inflammation is that it causes the body to produce too many harmful molecules called free radicals, while also reducing the levels of protective substances known as antioxidants. These changes can harm cells and lead to various diseases 5.

- ***Acute inflammation*** is the body's first reaction to sudden biological stress. Examples include bacterial infections or sudden tissue injuries. An inflammatory response is the body's way of reacting to injury or infection. When something harmful enters the body, like germs or a cut, the immune system sends signals to help fight it off. This can cause redness, swelling, heat, and pain in the affected area. The goal of this response is to protect the body and start the healing process. This answer includes a series of events that lead to... Events that involve many different types of cells and molecules that find harmful germs or injured tissue, then signal and gather help. Other cells and helper molecules work together to remove the problem. "Agents, and then bring the body back to balance. "Responses come with physical signs that show how we feel." "With fever, an increased heart rate, and changes in blood flow." "The movement to tissues, which helps improve the different" "Methods of protection that are used." The inflammation process begins when the body detects an injury or infection. This triggers the immune system to respond, sending special cells to the affected area to help fight off harmful germs and start healing. A strong response from the body can also harm healthy tissues. Causing more damage to the tissue *and increasing stimulation 6.*




Fig:1 Schematic overview of the modern inflammation process, illustrating the initiation, amplification, and resolution phases with key cellular and molecular mediators [7].

MEDIATORS

Mediators play a crucial role in starting the process of acute inflammation. Important substances like vasoactive amines, lipid-derived eicosanoids, cytokines, and chemokines work together to manage changes in blood vessels and help recruit inflammatory cells to the affected area. Cell-adhesion molecules are essential because they help inflammatory cells move from the bloodstream to where inflammation is happening. Pro-inflammatory cytokines, including tumor necrosis factor (TNF) and interleukin-1 beta (IL-1β), activate specific signaling pathways in the cells lining blood vessels (called endothelial cells). This activation leads to an increase in the production of adhesion molecules, which help capture circulating white blood cells, known as leukocytes, and direct them to the site of inflammation.[8]

International Journal of Pharmacy and Pharmaceutical Research (IJPPR)
Volume 32, Issue 7, July 2026 ijppr.humanjournals.com ISSN: 2349-7203




Fig:2. Overview of the Types of Mediators[9]

Leukotriene C4 (LT C4) and its related compounds, LTD4 and others, are important substances in the body that play a role in various biological processes. The substance that reacts slowly during anaphylaxis. Leukotrienes are chemicals that the body produces in response to inflammation or injury. They play a significant role in the immune system by helping to manage how the body reacts to allergens and infections. These substances can cause the airways to tighten, leading to breathing difficulties, especially in conditions like asthma. Leukotrienes are important for developing treatments for allergic reactions and respiratory issues. Leukotrienes are produced by almost all types of cells that are involved in Inflammatory responses, such as mast cells and basophils. Eosinophils, neutrophils, and monocytes are types of white blood cells. Leukotrienes are produced from arachidonic acid, which is a type of fatty acid.

Phospholipase A2 helps convert arachidonic acid through a process involving five steps. Lipoxygenase converts substances into 5S-hydroperoxy-6,8-trans-11,14-cis-. Eicosatetraenoic acid is a type of fatty acid that has twenty carbon atoms and four double bonds in its structure. It is important for various bodily functions and is found in certain types of fish and algae. The same pathway involving 5-lipoxygenase... The method speeds up the transformation of 5S-hydroperoxy-6,8- trans-11,14-cis-eicosatetraenoic acid into leukotriene A4 (LTA4). The number "15" represents a quantity that is greater than 14 and less than 16. It is often used in various contexts, such as counting, measuring, or identifying a specific group. LTA4 can be changed into LTB4.

HISTAMINE

Histamine is a substance that helps widen blood vessels. "Tightens smooth muscle and is a strong stimulant of Blood vessel openness, lung mucus, and stomach." Acid secretion refers to the process where the stomach produces acid to help digest food. This acid breaks down food and helps kill harmful bacteria. It impacts different types of cells. Types include smooth muscle cells, nerve cells, and gland cells. Endocrine and exocrine cells, blood cells, and other types of cells. The immune system is the body's defense against germs and diseases. It helps protect us by identifying and attacking harmful invaders like bacteria and viruses. When the immune system works well, it keeps us healthy and fights off infections. The text discusses how something is important not just for immediate needs. Histamine can activate H2 receptors during hypersensitivity reactions. Tor-mediated anti-inflammatory effects, including inhibition of... "release of enzymes from human neutrophils." Blocking the release of histamine caused by IgE. "Peripheral white blood cells and the activation of suppressor T-cells." A lymphocyte is a type of white blood cell that plays a key role in the immune system. It helps the body fight off infections and diseases by recognizing and attacking harmful germs and foreign substances. There are different kinds of lymphocytes.

PROSTAGLANDINS

Prostaglandins are the substances that help trigger allergic reactions**.** The activation of immune cells called mast cells in human lungs is PGD2 is a type of chemical that our body produces. They become active when triggered by the IgE receptor or by calcium. Prostaglandins are produced by an enzyme called cyclooxygenase. enzyme linked to the endoplasmic reticulum of Mast cells are a type of cell found in the body's tissues. The main types of prostaglandins are PGD2, PGE2, and PGFα.

PLATELET ACTIVATING FACTOR

Platelet activating factor (PAF) is an ether-linked phospholipid so designated because of its discovery as a basophil-derived mediator of rabbit platelet activation. PAF can be produced by a number of cells involved in the inflammatory response, including mast cells, macrophages, neutrophils, and eosinophils.PAF synthesis proceeds by a 2-step pathway in which phospholipase A2 hydrolyzes 1-O-alkyl-2-acyl-glyc-eryl-3-phosphorylcholine to yield 1-O-alkyl-2-acyl-glyceryl-3-phosphocholine (Liso-PAF), which is then converted by the enzyme acetyltransferase to PAF. The biological activities of PAF include platelet activation, neutrophil activation, and smooth muscle contraction. Because PAF is rapidly inactivated in the body, it is likely that it induces a cascade of inflammatory events that stimulates the accumulation of eosinophils on the endothelial surface, which may be the first step in the recruitment of eosinophils to tissues. Eosinophils are also a source of PAF and may attract additional eosinophils, which may increase the inflammatory response. PAF stimulates eosinophils to release basic proteins, leading to epithelial cell damage and increased expression of low-affinity IgE receptors on eosinophils and monocytes.

CYTOKINES AND CHEMOKINES

Mast cells are a source of many multifunctional cytokines. Cytokines such as IL-1, -2, -5, -6, TNF-α, and GM-CSF influence allergic inflammation by regulating the activity of inflammatory cells, including macrophages, T lymphocytes, B lymphocytes, and eosinophils. IL-1 enhances T helper cell development as well as B cell development and proliferation, while IL-2 induces T cell proliferation and B cell activation. IL-4 induces the differentiation of B cells into IgE-secreting plasma cells and, together with TNF-α, upregulates expression of high- and low-affinity IgE receptors on antigen-presenting cells. IL-5 induces the activation of B cells and increases the differentiation and longevity of eosinophils, and IL-6 increases the synthesis and secretion of B-cell immunoglobulins. In clinical studies in patients with seasonal allergic rhinitis, nasal biopsies showed that immunoreactivity for IL- and IL-6 was concentrated mainly in mast cells, whereas IL-5 was found in mast cells and eosinophils. Similar findings of increased levels of cytokines (IL- and IL-10) in nasal secretions were reported in a study of children with seasonal allergic rhinitis. Chemokines such as IL-8, macrophage inflammatory protein 1α, macrophage chemotactic protein 1, -2, and -3, and RANTES are cytokines that function primarily as chemoattractant molecules for macrophages and circulating leukocytes38, but also induce other inflammatory effects, including histamine release by basophils and mast cells and eosinophil activation acidosis. Increased levels of the chemokines IL-8, macrophage inflammatory protein 1α and RANTES, which correlated with total symptom scores, as well as the cytokines TNF-α and GM-CSF, were detected in nasal secretions. Following allergen challenge in patients with allergic rhinitis. Inflammatory changes occurring during the allergic response produce priming of nasal tissues. The resulting hypersensitivity of the nose leads to an increase in symptomatic and inflammatory responses to allergic irritation or further irritation of the nasal mucosa.[10]

KININ

Strong evidence shows that kinins are rapidly produced after tissue damage. They appear to regulate many of the processes involved in inflammation. These include the widening of blood vessels (vasodilation), increased permeability of blood vessel walls, leakage of plasma from blood vessels, migration of inflammatory cells to the affected area, and the development of pain and hyperalgesia. Hyperalgesia refers to an increased sensitivity to pain, resulting in an exaggerated pain response. It has also been shown that some of the effects of kinins may occur through mechanisms independent of specific receptors. These effects are thought to be associated with the direct release of histamine and 5-hydroxytryptamine (serotonin) from mast cells, which further contribute to the inflammatory response.[11]

Causes and physical effects "harmful results from inflammation. The inflammation can vary based on what causes it. The response serves a different purpose in the body. and harmful effects. Among the three Possible starting points include infections. Induced inflammation is linked to the "starting an immune response."

International Journal of Pharmacy and Pharmaceutical Research (IJPPR)
Volume 32, Issue 7, July 2026 ijppr.humanjournals.com ISSN: 2349-7203




Fig:3 Causes, and physiological and pathological outcomes, of inflammation[12]

ANTI-INFLAMMATORY

The term "anti-inflammatory" describes a chemical, therapeutic agent, or intervention's capacity to control the body's immunological and inflammatory responses to avoid, lessen, or suppress inflammation. Infection, tissue damage, toxins, or autoimmune reactions can all cause inflammation, which is a complicated biological process. While excessive or chronic inflammation can contribute to the onset and progression of many diseases, such as arthritis, cardiovascular disorders, diabetes, neurodegenerative diseases, inflammatory bowel disease, and some types of cancer, acute inflammation is a protective mechanism necessary for tissue repair and host defence. As a result, anti-inflammatory medicines are essential for preserving physiological homeostasis and controlling inflammatory diseases.[13]

Anti-inflammatory drugs work in a variety of cellular and molecular ways. Inhibition of pro-inflammatory cytokines, such as tumour necrosis factor-alpha (TNF-α), interleukin-1 beta (IL-1β), interleukin-6 (IL-6), and interleukin-8 (IL-8), which are important mediators in the initiation and amplification of inflammatory reactions, is one of the main mechanisms. Major inflammatory signalling pathways are also suppressed by these drugs, especially the cyclooxygenase (COX) and nuclear factor-kappa B (NF-κB) pathways. While inhibition of cyclooxygenase enzymes (COX-1 and COX-2) inhibits the manufacture of prostaglandins, which are powerful mediators of pain, fever, and inflammation, inhibition of NF-κB affects the transcription of many inflammatory genes.[14]

Both in vitro and in vivo experimental paradigms are frequently used to assess anti-inflammatory effectiveness. In vitro research usually evaluates the activation of signalling pathways, cytokine synthesis, cyclooxygenase activity, nitric oxide release, and suppression of inflammatory mediators in cultured immune cells like macrophages. To assess the effectiveness of anti-inflammatory drugs in physiological settings, in vivo models such as carrageenan-induced paw edema, cotton pellet granuloma, xylene-induced ear edema, and formalin-induced inflammation are widely used. The underlying mechanisms of action are further examined using molecular techniques such as Western blotting, quantitative real-time PCR (qRT-PCR), enzyme-linked immunosorbent assay (ELISA), immunohistochemistry, and transcriptome analysis.[15]

DIAGNOSIS

The diagnosis of infections and sepsis in a lab still depends on diagnostic tests. Diagnostic tests have been available since the 1970s or even earlier. Decisions made by doctors about the chances of a patient having an infection or sepsis are usually based on: "Are usually created with:" White blood cell counts, specifically for neutrophils. Myeloid-immature cells can be found in the bloodstream. "Blood (bandemia or 'left-shift')": Laboratory tests for substances that change during inflammation, like C-reactive protein, are important in diagnosing conditions. C-reactive protein (CRP) is a clinical sign, along with fever and rapid breathing. The cellular diagnostic measurements often cause issues. They are not very specific, especially for babies.[16]

Inflammation plays a crucial role in various health problems. Doctors often notice specific signs during their medical examinations. The five common signs of inflammation include redness, swelling, pain, loss of function, and sometimes fever. Each of these signs can feel unique to each person and may vary in strength or even be absent altogether. To identify inflammation, medical professionals look for markers in the blood, which are especially important in the early stages of diagnosis. Approximately fifty years ago, doctors

primarily focused on counting white blood cells and red blood cells. They also examined the red blood cell sedimentation rate in the lab, which helped in detecting inflammation. More recent tests have shown that acute-phase proteins can affect the results, making it harder to get a clear picture of inflammation. The charge on the surface of red blood cells can cause them to behave differently during inflammation, leading to changes in the sedimentation rate. However, because of the slow response times and the lack of detailed information from these tests, the significance of the erythrocyte sedimentation rate has declined over the years. Despite this, the white blood cell count remains a valuable indicator, even if it isn't the most precise measure. Since the times of ancient Greece and Rome, inflammation has been recognized as a key factor in many health issues. "Doctors have found signs during their examinations." The five common signs are redness, swelling, pain, and loss of function. "Function (function laesa) and eventually, fever (Calor)."These signs are quite personal and can be interpreted in different ways. "Weak or completely missing." Finding markers in the blood. Samples are very important in the beginning stages. Finding, describing, and treating inflammation. Around fifty years ago, doctors mainly checked the number of white blood cells and the red blood cells in the blood. The lab had the results for the red blood cell sedimentation rate. "Finding inflammation." The main idea of the recent test shows that acute-phase proteins disrupt negative results. The charge on the surface of red blood cells causes an increase in red blood cell activity. "Red blood cell sedimentation during times of inflammation." However, due to slow response time and a lack of clear details, the... The importance of the erythrocyte sedimentation rate has decreased since then. Sure! Please provide the text you would like me to paraphrase. The number of white blood cells, while not very precise, is still important.[17]

TREATMENT

conventional anti-inflammatory treatments of illnesses like IBD contain a group of standard therapeutic treatments, to be specific, 5-ASAs, antimicrobials, steroids, immunosuppressants, immunomodulators, and biologics. Their essential capacities are symptom control, reduction, restraint of inflammation, and modulation of dysregulated immune reactions. Comparatively, the components of both probiotics and conventional treatments coincide with the common denominator of balancing inflammation and safety. Conventional drugs tend to do this by straightforwardly focusing on key molecular components or pathways inside inflammation and susceptibility, e.g., hindering the pro-inflammatory cytokines TNF-α, IL-6, and IL-1β, or balancing the major signaling transduction cascades JAK/STAT and NF-kB. Similarly, probiotics also have anti-inflammatory impacts through coordinated tweaking of these pathways by hindering NF-kB and JAK/STAT cascades and pro-inflammatory cytokines like IL-6 and IL-1β generation.[18]




Fig. (3). Mechanism of action of different compounds to treat inflammation. Inhibition of NF-κβ pathway. A schematic illustration of the steps involved in the activation of the NF-κβ pathway. Numerous drugs, natural products and recombinant compounds can act at several of these steps to interfere with NF-κβ activation.

Fig:4 Mechanism of action of different compounds to treat inflammation

International Journal of Pharmacy and Pharmaceutical Research (IJPPR)
Volume 32, Issue 7, July 2026 ijppr.humanjournals.com ISSN: 2349-7203




Fig:5. Summarized anti-inflammatory role of drugs and natural compounds in diseases.

1 Non-Steroidal Anti-Inflammatory Drugs (NSAIDs)

Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) speaks to a critical bunch of drugs with critical restorative applications to remedy fiery conditions such as rheumatoid joint pain or osteoarthritis [106,107], and have given a momentous advancement, beginning from the essential sedate headache medicine to the most recent rise and drop of particular COX-2 inhibitors.

2 Glucocorticoids

Glucocorticoids are a gather of steroid hormones discharged by the adrenal cortex after actuation of hypothalamus pituitary- adrenal pivot (HPA) by incendiary stretch related jolts, and their work is by and large interceded by means of cytosolic receptors, especially glucocorticoid receptors (GRs) that are related with the atomic hormone receptor super family and are ligand-induced translation components with an capacity to control quality expression by coordinate interaction with specific promoter arrangements.

3 PPAR- (Peroxisome Proliferator Activated Receptors) Agonists

PPAR- applies its anti-inflammatory exercises in both the fringe as well as the CNS glial and neuronal cells, where it controls the expression of pro-inflammatory particles iNOs, COX-2 and in the long run an broad bunch of cytokines through its intelligent with the translation calculate NF-B, taking after actuation by its actually happening ligands eicosanoids and PGJ2 the pioglitazone has moreover been as of late watched to display anti-inflammatory exercises conjointly reduce diabetic nephropathy by restraining NF-B enactment.

4 ACE (Angiotensinogen Converting Enzyme) Inhibitors

the treatment with Expert inhibitors seems invert the expanded generation of superoxide anions, NF- enactment, expression of pro-inflammatory cytokines and diminished NO union in atherosclerosis. As of late, Captopril, a Pro inhibitor, has too been watched to operate as an anti-inflammatory operator in hypertensive creatures [99], and its anti-inflammatory movement has already been appeared in a few other illnesses such as colitis and joint pain that are characterized by an aggravation handle.[19]

Conclusion

An intricate and vital biological process, inflammation is vital to the body's Défense against injuries, infections, and damaging stimuli. Uncontrolled or prolonged inflammation can contribute to the onset and progression of a few chronic diseases, including neurodegenerative conditions, autoimmune diseases, metabolic disorders, and cardiovascular disorders, even though acute inflammation is a helpful Défense mechanism that encourages healing and tissue repair. Chronic inflammation, especially age-related inflammation or "inflammaging," is strongly connected with cellular stress, mitochondrial dysfunction, DNA damage, and cellular senescence, all of which hasten the course of disease.

Numerous cellular and molecular mediators, including cytokines, chemokines, eicosanoids, and adhesion molecules, control inflammatory responses. Knowing these pathways has made it easier to identify diagnostic markers like C-reactive protein. Numerous cellular and molecular mediators, including cytokines, chemokines, eicosanoids, and adhesion molecules, control inflammatory responses. Knowing these pathways has made it easier to identify diagnostic indicators that aid in the early detection and monitoring of inflammatory disorders, such as blood cell alterations, inflammatory cytokines, and C-reactive protein (CRP). However, the complexity and variety of inflammatory responses continue to make correct diagnosis difficult. Restoring physiological homeostasis and reducing excessive immune activation are important goals of anti-inflammatory treatments.

Non-steroidal anti-inflammatory medications (NSAIDs), glucocorticoids, PPAR agonists, ACE inhibitors, immunosuppressants, and biologic therapies are examples of conventional treatments that work by focusing on important inflammatory pathways such as NF-κB, COX enzymes, and pro-inflammatory cytokines. The promise of more recent methods, such as targeted molecular medicines and biological interventions, to offer safer and more efficient treatment alternatives has also been emphasized by advances in our understanding of inflammation.

In general, inflammation is a two-edged process: while it is necessary for healthy Défense and repair, it may also be detrimental if it persists or is dysregulated. Developing better methods to prevent and treat inflammation-related illnesses while preserving immune system balance requires ongoing study into inflammatory causes, biomarkers, and therapeutic approaches.

REFERENCES

1 Xiao L, Harrison DG. Inflammation in hypertension. Canadian journal of cardiology. 2020 May 1;36(5):635-47
2 Gupta L, Thomas J, Ravichandran R, Singh M, Nag A, Panjiyar BK. Inflammation in cardiovascular disease: A comprehensive review of biomarkers and therapeutic targets. Cureus. 2023 Sep 18;15(9).
3 Pezone A, Olivieri F, Napoli MV, Procopio A, Avvedimento EV, Gabrielli A. Inflammation and DNA damage: cause, effect, or both. Nature Reviews Rheumatology. 2023 Apr;19(4):200-11.
4 Medzhitov R. Origin and physiological roles of inflammation. Nature. 2008 Jul 24;454(7203):428-35
5 Khansari N, Shakiba Y, Mahmoudi M. Chronic inflammation and oxidative stress as a major cause of age-related diseases and cancer. Recent patents on inflammation & allergy drug discovery. 2009 Jan 1;3(1):73-80.
6 Vodovotz Y, Chow CC, Bartels J, Lagoa C, Prince JM, Levy RM, Kumar R, Day J, Rubin J, Constantine G, Billiar TR. In silico models of acute inflammation in animals. Shock. 2006 Sep 1;26(3):235-44
7 . Egamberdieva G. INFLAMMATION AS A THERAPEUTIC TARGET IN CHRONIC INTERNAL DISEASES: EMERGING EVIDENCE AND CLINICAL IMPLICATIONS. Journal of Clinical and Biomedical Research. 2026 Jan 7;1(1):28-34.
8 Lawrence T, Willoughby DA, Gilroy DW. Anti-inflammatory lipid mediators and insights into the resolution of inflammation. Nature Reviews Immunology. 2002 Oct 1;2(10):787-95.
9 Ansar W, Ghosh S. Inflammation and inflammatory diseases, markers, and mediators: role of CRP in some inflammatory diseases. InBiology of C reactive protein in health and disease 2016 Mar 24 (pp. 67-107). New Delhi: Springer India.
10 White M. Mediators of inflammation and the inflammatory process. Journal of Allergy and Clinical Immunology. 1999 Mar 1;103(3): S378-81.
11 Calixto JB, Cabrini DA, Ferreira J, Campos MM. Kinins in pain and inflammation. Pain. 2000 Jul 1;87(1):1
12 Medzhitov R. Origin and physiological roles of inflammation. Nature. 2008 Jul 24;454(7203):428-35.
13 Robbin & Cotran Pathologic Basis of Disease Kumar, V., Abbas, A. K., & Aster, J. C. (2020). Robbins & Cortan Pathologic Basis of Disease (10th ed.). Elsevier
14 Rang & Dale's Pharmacology Ritter, J.M., Flower, R.J., Henderson, G., Loke, Y. k., & MacEwan, D. (2023). Rang & Dale's Pharmacology (10th ed.). Elsevier
15 Ruslan Medzhitov. (2008). Origin and physiological roles of inflammation. Nature, 454(7203), 428-435. A highly cited review discussing the biology of inflammation and inflammatory mediators.
16 Davis BH. Improved diagnostic approaches to infection/sepsis detection. Expert review of molecular diagnostics. 2005 Mar 1;5(2):193-207.

17 Pfäfflin A, Schleicher E. Inflammation markers in point-of-care testing (POCT). Analytical and bioanalytical chemistry. 2009 Mar;393(5):1473-80.
18 Beikmohammadi M, Halimi S, Fard NA, Wen W. Therapeutic potential of probiotics: A review of their role in modulating inflammation. Probiotics and Antimicrobial Proteins. 2026 Jan;18(1):1122-41
19 Laveti D, Kumar M, Hemalatha R, Sistla R, Gm Naidu V, Talla V, Verma V, Kaur N, Nagpal R. Anti-inflammatory treatments for chronic diseases: a review. Inflammation & Allergy-Drug Targets-Inflammation & Allergy). 2013 Oct 1;12(5):349-61.

How to cite this article:
Mr. Shantakumar et al. Ijppr.Human, 2026; Vol. 32 (7):390-398.
Conflict of Interest Statement: All authors have nothing else to disclose.
This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.